

Department of Mathematics, Indian Institute of Technology, New Delhi, India

Department of Mathematics, Indian Institute of Technology, Kharagpur, India

N. K. GOVIL and V. K. JAIN

Some Integral Inequalities for Entire Functions of Exponential Type

Pewne nierówności całkowe dla funkcji całkowitych typu wykładniczego

Некоторые интегральные неравенства для целых функций экспоненциального типа

1. INTRODUCTION AND STATEMENT OF RESULTS

Let $f(z)$ be an entire function of exponential type τ . The following integral inequalities (for references, see [1, pp. 211, 98]) are well known.

THEOREM A. If $f(z)$ is an entire function of exponential type τ belonging to L^p ($1 \leq p < \infty$) on the real axis, then

$$(1.1) \quad \int_{-\infty}^{\infty} |f'(x)|^p dx \leq \tau^p \int_{-\infty}^{\infty} |f(x)|^p dx$$

and

$$(1.2) \quad \int_{-\infty}^{\infty} |f(x + iy)|^p dx \leq e^{p\tau|y|} \int_{-\infty}^{\infty} |f(x)|^p dx, \quad -\infty < y < \infty.$$

If $h_f(\pi/2) = 0$ ($h_f(\theta) = \limsup_{r \rightarrow \infty} \frac{\log |f(re^{i\theta})|}{r}$ is the indicator function of $f(z)$) and $f(z) \neq 0$ for $\text{Im } z > 0$, then the inequality analogous to (1.1) has been obtained by Rahman [5]. No inequality analogous to (1.2) is known, but if $p = 2$, it has been proved by Rahman [6] that for $y < 0$,

$$(1.3) \quad \int_{-\infty}^{\infty} |f(x + iy)|^2 dx \leq \frac{e^{2\tau|y|} + 1}{2} \int_{-\infty}^{\infty} |f(x)|^2 dx .$$

An inequality analogous to (1.1) for functions of exponential type not vanishing in $\text{Im } z > k$ ($k \leq 0$) has been obtained by Govil and Rahman [2]. In this paper we consider the class of entire functions of exponential type τ satisfying $f(z) \equiv \omega(z)$, where $\omega(z) = e^{i\tau z} \overline{f(z)}$ and prove the following

THEOREM 1. Let $f(z)$ be an entire function of exponential type τ belonging to L^p ($1 \leq p < \infty$) on the real axis. If $f(z) \equiv \omega(z)$, then we have

$$(1.4) \quad \int_{-\infty}^{\infty} |f'(x)|^p dx \leq C_p \tau^p \int_{-\infty}^{\infty} |f(x)|^p dx$$

and

$$(1.5) \quad \int_{-\infty}^{\infty} |f'(x)|^p dx \geq (1 - C_p^p)^p \tau^p \int_{-\infty}^{\infty} |f(x)|^p dx ,$$

where

$$C_p = \frac{2\pi}{\int_0^{2\pi} |1 + e^{i\alpha}|^p d\alpha}$$

THEOREM 2. If $f(z)$ is an entire function of exponential type τ belonging to L^2 on the real axis and satisfying $f(z) \equiv \omega(z)$, then

$$(1.6) \quad \int_{-\infty}^{\infty} |f(x + iy)|^2 dx \leq \frac{e^{-2\tau y} + 1}{2} \int_{-\infty}^{\infty} |f(x)|^2 dx, \quad -\infty < y < \infty$$

THEOREM 3. Let $f(z)$ be an entire function of exponential type τ (≥ 1) and periodic on the real axis with period 2π . If $f(z) = \omega(z)$, then

$$(1.7) \quad \int_{-\pi}^{\pi} |f'(x)|^2 dx \leq \frac{\tau^2}{2} \int_{-\pi}^{\pi} |f(x)|^2 dx$$

and

$$(1.8) \quad \int_{-\pi}^{\pi} |f(x + iy)|^2 dx \leq \frac{e^{-2\tau y} + 1}{2} \int_{-\pi}^{\pi} |f(x)|^2 dx, \quad -\infty < y < \infty$$

We also prove

THEOREM 4. Let $f(z)$ be an entire function of exponential type τ (≥ 1) and periodic on the real axis with period 2π . If $f(z) = e^{i\tau z} f(-z)$, then

$$(1.9) \quad \int_{-\pi}^{\pi} |f'(x)|^2 dx \leq \frac{\tau^2}{2} \int_{-\pi}^{\pi} |f(x)|^2 dx$$

and

$$(1.10) \quad \int_{-\pi}^{\pi} |f(x + iy)|^2 dx \leq \frac{e^{-2\tau y} + 1}{2} \int_{-\pi}^{\pi} |f(x)|^2 dx, \quad -\infty < y < \infty$$

2. LEMMAS

LEMMA 1. If $f(z)$ is regular and of exponential type in the upper half plane, $h_p(\pi/2) \leq c$ and $|f(x)| \leq M$, $-\infty < x < \infty$ then

$$(2.1) \quad |f(x + iy)| \leq M e^{cy}, \quad -\infty < x < \infty, \quad 0 \leq y < \infty$$

This Lemma is due to Pólya and Szegő [4, p. 36], Boas [1, p. 82].

LEMMA 2. If $f(z)$ is an entire function of exponential type τ belonging to L^p ($1 \leq p < \infty$) on the real axis, then

$$(2.2) \quad \int_{-\infty}^{\infty} |i\tau f(x) + f'(x) + e^{i\alpha} \{-i\tau f(x) + f'(x)\}|^p dx \\ \leq (2\tau)^p \int_{-\infty}^{\infty} |f(x)|^p dx, \quad \alpha \in [0, 2\pi).$$

This Lemma is due to Rahman [5, inequality (3.18), p. 300]

3. PROOFS OF THEOREMS

P r o o f of Theorem 1. Since $f(z)$ is an entire function of exponential type τ , belonging to L^p ($1 \leq p < \infty$) on real axis, there exists a constant M [Boas [1, Th. 6.7.1]] such that $|f(x)| \leq M$, $-\infty < x < \infty$. Further since $f(z) = \omega(z)$ we have $h_p(\pi/2) \leq 0$. Now if $f_1(z)$ denotes the function $e^{-i\tau z/2} f(z)$, then $f_1(z)$ is of exponential type $\tau/2$ and belongs to L^p ($1 \leq p < \infty$). Hence applying Lemma 2 to $f_1(z)$, we get

$$\int_{-\infty}^{\infty} \left| 1 - \frac{\tau}{2} f_1(x) + f_1'(x) + e^{i\alpha} \left\{ -1 - \frac{\tau}{2} f_1(x) + f_1'(x) \right\} \right|^p dx$$

$$\leq \tau^p \int_{-\infty}^{\infty} |f_1(x)|^p dx, \quad (p \geq 1)$$

which gives

$$\int_{-\infty}^{\infty} \left| f'(x) e^{-i\tau x/2} + e^{i\alpha} \left\{ -i\tau e^{-i\tau x/2} f(x) + e^{-i\tau x/2} f'(x) \right\} \right|^p dx \leq \tau^p \int_{-\infty}^{\infty} |f(x)|^p dx \quad (p \geq 1).$$

Consequently

$$\int_{-\infty}^{\infty} \left| f'(x) + e^{i\alpha} \left\{ -i\tau f(x) + f'(x) \right\} \right|^p dx \leq \tau^p \int_{-\infty}^{\infty} |f(x)|^p dx, \quad (p \geq 1).$$

Integrating both sides with respect to α from 0 to 2π , we get

$$(3.1) \quad \int_0^{2\pi} d\alpha \int_{-\infty}^{\infty} \left| f'(x) + e^{i\alpha} \left\{ -i\tau f(x) + f'(x) \right\} \right|^p dx$$

$$\leq 2\pi \tau^p \int_{-\infty}^{\infty} |f(x)|^p dx \quad (p \geq 1).$$

Note that $f'(x)$ can be zero only at a countable number of points. Besides, we can clearly invert the order of integration on the left side of (3.1). Therefore

$$(3.2) \quad \int_0^{2\pi} d\alpha \int_{-\infty}^{\infty} \left| f'(x) + e^{i\alpha} \left\{ -i\tau f(x) + f'(x) \right\} \right|^p dx =$$

$$= \int_0^{2\pi} d\alpha \int_{-\infty}^{\infty} |f'(x)|^p \left| 1 + e^{i\alpha} \frac{-i\tau f(x) + f'(x)}{f'(x)} \right|^p dx =$$

$$\begin{aligned}
 &= \int_{-\infty}^{\infty} |f'(x)|^p dx \int_0^{2\pi} \left| 1 + e^{i\alpha} \frac{-i\tau f(x) + f'(x)}{f'(x)} \right|^p d\alpha = \\
 &= \int_{-\infty}^{\infty} |f'(x)|^p dx \int_0^{2\pi} \left| 1 + e^{i\alpha} \frac{B(x)}{A(x)} \right|^p d\alpha,
 \end{aligned}$$

where $B(x) = -i\tau f(x) + f'(x)$ and $A(x) = f'(x)$.

Further since $f(z) \equiv \omega(z)$, we have for real x

$$\begin{aligned}
 |A(x)| &= |f'(x)| \\
 &= |\omega'(x)| \\
 &= |-i\tau f(x) + f'(x)| \\
 &= |B(x)|,
 \end{aligned}$$

i.e. $\left| \frac{B(x)}{A(x)} \right| = 1$. Thus for a fixed real x and every $p > 0$

$$(3.3) \quad \int_0^{2\pi} \left| 1 + e^{i\alpha} \frac{B(x)}{A(x)} \right|^p d\alpha = \int_0^{2\pi} |1 + e^{i\alpha}|^p d\alpha, \quad (p > 0)$$

Combining inequality (3.1) and equalities (3.2), (3.3), we get

$$\int_0^{2\pi} |1 + e^{i\alpha}|^p d\alpha \int_{-\infty}^{\infty} |f'(x)|^p dx \leq 2\pi \tau^p \int_{-\infty}^{\infty} |f(x)|^p dx, \quad (p \geq 1)$$

which is (1.4).

To prove (1.5), note that $\omega(z)$ is an entire function of exponential type τ satisfying $\omega(z) = e^{i\tau z} \{\overline{\omega(\bar{z})}\}$.

Hence using (1.4), we get

$$\int_{-\infty}^{\infty} |\omega'(x)|^p dx \leq C_p \tau^p \int_{-\infty}^{\infty} |\omega(x)|^p dx,$$

which is equivalent to

$$(3.4) \quad \left(\int_{-\infty}^{\infty} |-1\tau f(x) + f'(x)|^p dx \right)^{\frac{1}{p}} \leq \tau C_p \left(\int_{-\infty}^{\infty} |f(x)|^p dx \right)^{\frac{1}{p}},$$

($p \geq 1$).

Therefore by Minkowski's inequality

$$\begin{aligned} & \left(\int_{-\infty}^{\infty} \{ |-1\tau f(x) + f'(x)| + |-f'(x)| \}^p dx \right)^{\frac{1}{p}} \\ & \leq \left(\int_{-\infty}^{\infty} \{ |-1\tau f(x) + f'(x)| \}^p dx \right)^{\frac{1}{p}} + \left(\int_{-\infty}^{\infty} |f'(x)|^p dx \right)^{\frac{1}{p}}, \end{aligned}$$

which gives

$$(3.5) \quad \tau \left(\int_{-\infty}^{\infty} |f(x)|^p dx \right)^{\frac{1}{p}} \leq \left(\int_{-\infty}^{\infty} |-1\tau f(x) + f'(x)|^p dx \right)^{\frac{1}{p}} + \left(\int_{-\infty}^{\infty} |f'(x)|^p dx \right)^{\frac{1}{p}}.$$

On combining (3.4) and (3.5), we get

$$\tau \left(1 - C_p^p \right) \left(\int_{-\infty}^{\infty} |f(x)|^p dx \right)^{\frac{1}{p}} \leq \left(\int_{-\infty}^{\infty} |f'(x)|^p dx \right)^{\frac{1}{p}}.$$

from which (1.5) follows.

P r o o f of Theorem 2. Since $f(z) \in L^2$ on the real axis, we have by Paley-Wiener Theorem [3, pp. 499-501]

$$(3.6) \quad f(z) = \int_0^{\tau} e^{1zt} \varphi(t) dt, \quad \varphi \in L^2(0, \tau).$$

Now

$$(3.7) \quad \omega(z) = e^{1\tau z} \int_0^{\tau} e^{-1zt} \overline{\varphi(t)} dt$$

$$= \int_0^{\tau} e^{iz(\tau-t)} \overline{\varphi(t)} dt$$

Since $f(z) \equiv \omega(z)$, hence

$$\begin{aligned} \int_{-\infty}^{\infty} |f(x+iy)|^2 dx &= \frac{1}{2} \int_{-\infty}^{\infty} |\omega(x+iy)|^2 dx + \frac{1}{2} \int_{-\infty}^{\infty} |f(x+iy)|^2 dx \\ &= \pi \int_0^{\tau} e^{-2y(\tau-t)} |\varphi(t)|^2 dt + \pi \int_0^{\tau} e^{-2yt} |\varphi(t)|^2 dt \\ &\leq \pi(e^{-2\tau y} + 1) \int_0^{\tau} |\varphi(t)|^2 dt \\ &= \frac{(e^{-2\tau y} + 1)}{2} \int_{-\infty}^{\infty} |f(x)|^2 dx, \end{aligned}$$

which is (1.6).

P r o o f of Theorem 3. Since $f(z)$ is an entire function of exponential type τ and is periodic on the real axis with period 2π , we have (see Boas [1, p. 109])

$$(3.8) \quad f(z) = \sum_{k=-n}^n a_k e^{ikz}, \quad n \leq \tau$$

and since $f(z) \equiv \omega(z)$, we have $h_p(\pi/2) \leq 0$. Hence we get from (3.8)

$$(3.9) \quad f(z) = \sum_{k=0}^n a_k e^{ikz}, \quad n \leq \tau.$$

Further

$$(3.10) \quad \omega(z) = e^{i\tau z} \sum_{k=0}^n \overline{a_k} e^{-ikz} = \sum_{k=0}^n \overline{a_k} e^{i(\tau-k)z}$$

Therefore

$$\begin{aligned} \int_{-\pi}^{\pi} |f'(x)|^2 dx &= \frac{1}{2} \int_{-\pi}^{\pi} |f'(x)|^2 dx + \frac{1}{2} \int_{-\pi}^{\pi} |\omega'(x)|^2 dx = \\ &= \pi \sum_{k=0}^n k^2 |a_k|^2 + \pi \sum_{k=0}^n (\tau - k)^2 |a_k|^2, \end{aligned}$$

(by (3.9) and (3.10))

$$\leq \pi \tau^2 \sum_{k=0}^n |a_k|^2 = \frac{\tau^2}{2} \int_{-\pi}^{\pi} |f(x)|^2 dx,$$

which is (1.7).

To prove (1.8), we have

$$\begin{aligned} \int_{-\pi}^{\pi} |f(x + iy)|^2 dx &= \frac{1}{2} \int_{-\pi}^{\pi} |f(x + iy)|^2 dx + \frac{1}{2} \int_{-\pi}^{\pi} |\omega(x + iy)|^2 dx = \\ &= \pi \sum_{k=0}^n e^{-2ky} |a_k|^2 + \pi \sum_{k=0}^n e^{-2(\tau-k)y} |a_k|^2 \\ &\leq \pi (1 + e^{-2\tau y}) \sum_{k=0}^n |a_k|^2 = \\ &= \frac{1 + e^{-2\tau y}}{2} \int_{-\pi}^{\pi} |f(x)|^2 dx, \end{aligned}$$

which is (1.8).

P r o o f of Theorem 4. Here $f(z) = e^{i\tau z} f(-z)$. Hence we get $h_f(\pi/2) \leq 0$. And so here also, the representation (3.8) of $f(z)$ will reduce to the representation (3.9).

Then the proof follows on the lines similar to that of Theorem

3.

REFERENCES

- [1] Boas, R.P., Jr., Entire functions, Academic Press, New York 1954.
- [2] Govil, N.K., Rahman, Q.I., Functions of exponential type not vanishing in a half plane and related polynomials, Trans. Amer. Math. Soc., 137(1969), 501-517.
- [3] Levin, B.Ya., Distribution of zeros of entire functions (Russian), Moscow 1956.
- [4] Pólya, G., Szegő, G., Aufgaben and Lehrsätze aus der Analysis, Vol. II, Springer, Berlin, 1925.
- [5] Rahman, Q.I., Functions of exponential type, Trans. Amer. Math. Soc., 135(1969), 295-309.
- [6] ,, ,, Inequalities for polynomials and entire functions, Illinois J. Math., 5(1961), 141-151.

STRESZCZENIE

W pracy udowodniono nierówność (1)

$$(1) \quad \int_{-\infty}^{+\infty} |f'(x)|^p dx \leq c_p \tau^p \int_{-\infty}^{+\infty} |f(x)|^p dx$$

dla funkcji całkowitej $f \in L^p$, $p \geq 1$ typu wykładniczego τ oraz przy warunku, że $f(z) = e^{1/\tau z} \overline{f(\bar{z})}$, nierówność (2)

$$(2) \quad \int_{-\infty}^{+\infty} |f'(x)|^2 dx \leq \frac{e^{-2\tau y} + 1}{2} \int_{-\infty}^{+\infty} |f(x)|^2 dx, \quad y < 0.$$

Ponadto otrzymano nierówność przeciwną do (1) z zamianą c_p na $(1 - c_p^p)^{1/p}$ oraz kilka innych analogicznych nierówności do (1) i (2).

Резюме

В работе доказано неравенство $(1) \int_{-\infty}^{+\infty} |f'(x)|^p dx \leq c_p \tau^p \int_{-\infty}^{+\infty} |f(x)|^p dx$ для целой функции $f \in L^p$, $p \geq 1$ экспоненциального типа τ , а также, при условии $f(z) = e^{i\tau z} \overline{f(\bar{z})}$, неравенство (2)

$$(2) \int_{-\infty}^{+\infty} |f'(x)|^2 dx \leq \frac{e^{-2\tau y} + 1}{2} \int_{-\infty}^{+\infty} |f(x)|^2 dx, \quad y < 0.$$

Кроме того получено неравенство противоположное к (1) с заменой c_p на $(1 - c_p^{\frac{1}{p}})^p$, а также несколько других аналогических неравенств к (1) и (2).

