

FURKAN SEÇGIN

## On Leonardo, Leonardo–Lucas and modified Leonardo elliptic quaternions and their matrix representations

**ABSTRACT.** In this paper, we present a new class of elliptic quaternions that incorporate Leonardo, Leonardo–Lucas and modified Leonardo numbers into their components. We explore some fundamental properties associated with these numbers. In particular, we obtain recurrence relations, generating function, Binet formula of these sequences and by using Binet formula we derive Vajda, Cassini, Catalan and d’Ocagne identities. Lastly, we investigate two different matrix representations of these numbers.

**1. Introduction.** Recently, Catarino and Borges [3] explored the recurrence relations and various properties of Leonardo numbers. Following their work, Alp and Koçer [1] investigated additional interesting properties of these numbers. Kuhapatanakul and Juthamas [8] extended the study to generalized Leonardo numbers, examining their matrix representation. Additionally, Karatas [7] defined complex Leonardo numbers and analyzed their combinatorial properties. İşbilir et al. [6] investigated Pauli–Leonardo quaternions.

Further developments on Leonardo numbers, their generalizations, and interesting properties can be found in [2, 9, 12, 11, 14, 16, 17].

---

2010 *Mathematics Subject Classification.* 15B33, 40B05, 15A24, 11B83.

*Key words and phrases.* Quaternions, elliptic quaternions, Leonardo numbers, matrix representation.

Özdemir [10] explained how, for a given ellipsoid, one can define a suitable scalar product and vector product. He introduced elliptical inner product, elliptical vector product and he aimed to describe the motion on an ellipsoid  $x^2/a^2 + y^2/b^2 + z^2/c^2 = 1$  as a rotation using the proper inner product, vector and elliptical orthogonal matrices introduced in that elliptical scalar product space. He also defined the elliptic quaternions and showed a relation between elliptical rotation and elliptic quaternions.

Let  $\{e_0, e_1, e_2, e_3\}$  be four basic elements satisfying the equalities  $e_0 = 1$ ,  $e_1^2 = -\alpha$ ,  $e_2^2 = -\beta$ ,  $e_3^2 = -\gamma$  and

$$(1.1) \quad e_1e_2 = \frac{\Delta}{\gamma}e_3 = -e_2e_1, \quad e_2e_3 = \frac{\Delta}{\alpha}e_1 = -e_3e_2, \quad e_3e_1 = \frac{\Delta}{\beta}e_2 = -e_1e_3$$

where  $\alpha, \beta, \gamma \in \mathbb{R}^+$  and  $\Delta = \sqrt{\alpha\beta\gamma}$ . The set of elliptic quaternions is defined by the formula

$$(1.2) \quad \mathbb{H}_{\alpha,\beta,\gamma} = \{Q = q_0e_0 + q_1e_1 + q_2e_2 + q_3e_3 : q_0, q_1, q_2, q_3 \in \mathbb{R}\}.$$

This set is an associative, noncommutative division algebra with the basic elements  $\{e_0, e_1, e_2, e_3\}$  and it is a 4-dimensional algebra over real numbers. If  $\alpha, \beta$  and  $\gamma$  are equal to 1, we have the real quaternion algebra. Thus, elliptic quaternions are an extension of real quaternions with the elliptic inner and vector product. Furthermore, the following table shows the products of dual elliptic quaternion units.

$\cdot$	$e_0$	$e_1$	$e_2$	$e_3$
$e_0$	$e_0$	$e_1$	$e_2$	$e_3$
$e_1$	$e_1$	$-\alpha$	$(\Delta/\gamma)e_3$	$(-\Delta/\beta)e_2$
$e_2$	$e_2$	$(-\Delta/\gamma)e_3$	$-\beta$	$(\Delta/\alpha)e_1$
$e_3$	$e_3$	$(\Delta/\beta)e_2$	$(-\Delta/\alpha)e_1$	$-\gamma$

TABLE 1. The products of elliptic quaternionic units.

Additionally, as in the case of quaternions, it is possible to define right and left multiplications of two elliptic quaterions using  $4 \times 4$  real matrices. For an elliptic quaternion  $Q = q_0 + q_1e_1 + q_2e_2 + q_3e_3$ , consider the transformations

$$\varphi : \mathbb{H}_{\alpha,\beta,\gamma} \longrightarrow \mathbb{H}_{\alpha,\beta,\gamma} \quad \text{and} \quad \mu : \mathbb{H}_{\alpha,\beta,\gamma} \longrightarrow \mathbb{H}_{\alpha,\beta,\gamma}$$

$$P \longrightarrow \varphi(P) = QP \quad \text{and} \quad P \longrightarrow \mu(P) = PQ.$$

The following  $4 \times 4$  matrices corresponding to these transformations, respectively, provide the left and right product matrices of  $Q$ :

$$\mathcal{L}_Q = \begin{bmatrix} q_0 & -\alpha q_1 & -\beta q_2 & -\gamma q_3 \\ q_1 & q_0 & -\frac{\Delta}{\alpha} q_3 & \frac{\Delta}{\alpha} q_2 \\ q_2 & \frac{\Delta}{\beta} q_3 & q_0 & -\frac{\Delta}{\beta} q_1 \\ q_3 & -\frac{\Delta}{\gamma} q_2 & \frac{\Delta}{\gamma} q_1 & q_0 \end{bmatrix}, \quad \mathcal{R}_Q = \begin{bmatrix} q_0 & -\alpha q_1 & -\beta q_2 & -\gamma q_3 \\ q_1 & q_0 & \frac{\Delta}{\alpha} q_3 & -\frac{\Delta}{\alpha} q_2 \\ q_2 & -\frac{\Delta}{\beta} q_3 & q_0 & \frac{\Delta}{\beta} q_1 \\ q_3 & \frac{\Delta}{\gamma} q_2 & -\frac{\Delta}{\gamma} q_1 & q_0 \end{bmatrix}.$$

For detailed information about these concepts, we refer to reader to [4, 10, 13].

**2. Basic concepts and notions.** This section contains some definitions and theorems that will be used in the following sections of the paper.

For  $n \geq 2$ , the sequence of Fibonacci numbers  $\{F_n\}$  is defined by

$$(2.1) \quad F_n = F_{n-1} + F_{n-2}, \quad F_0 = 0, F_1 = 1$$

and the sequence of Lucas numbers  $\{L_n\}$  is defined by

$$(2.2) \quad L_n = L_{n-1} + L_{n-2}, \quad L_0 = 2, L_1 = 1.$$

Let  $\psi$  and  $\phi$  be the roots of the characteristic equation  $t^2 - t - 1 = 0$  corresponding to formulas (2.1)–(2.2). Solving the characteristic equation, we obtain the distinct roots  $\psi = \frac{1+\sqrt{5}}{2}$  and  $\phi = \frac{1-\sqrt{5}}{2}$ . Then the Binet formulas for Fibonacci ( $F_n$ ) and Lucas ( $L_n$ ) numbers are:

$$F_n = \frac{\psi^n - \phi^n}{\psi - \phi} \quad \text{and} \quad L_n = \psi^n + \phi^n.$$

Note that we have the following equations:

$$(2.3) \quad 2F_{n+1} = F_n + L_n, \quad L_n = F_{n-1} + F_{n+1}, \quad L_n = F_{n+2} - F_{n-2}.$$

Similar to the Fibonacci and Lucas numbers, the Leonardo numbers, which were first introduced by Dijkstra [5] in 1981 in a sorting algorithm, satisfy the recurrence relation

$$(2.4) \quad \mathcal{L}_n = \mathcal{L}_{n-1} + \mathcal{L}_{n-2} + 1, \quad n \geq 2$$

with the initial conditions  $\mathcal{L}_0 = \mathcal{L}_1 = 1$ . This sequence can be seen in OEIS [A001595]. Furthermore, there are two number sequences called Leonardo–Lucas and modified Leonardo introduced by Soykan in [14]. For  $n \geq 2$ , they are defined as

$$(2.5) \quad \mathcal{L}_{L,n} = \mathcal{L}_{L,n-1} + \mathcal{L}_{L,n-2} - 1, \quad \mathcal{L}_{L,0} = 3, \mathcal{L}_{L,1} = 2$$

and

$$(2.6) \quad \mathcal{M}_n = \mathcal{M}_{n-1} + \mathcal{M}_{n-2} + 1, \quad \mathcal{M}_0 = 0, \mathcal{M}_1 = 1,$$

respectively. These sequences can be seen in OEIS [A001612] and [A000071], respectively. The sequences  $\{\mathcal{L}_n\}$ ,  $\{\mathcal{L}_{L,n}\}$  and  $\{\mathcal{M}_n\}$  satisfy the third order linear recurrences as follows:

$$(2.7) \quad \begin{aligned} \mathcal{L}_n &= 2\mathcal{L}_{n-1} - \mathcal{L}_{n-3}, & \mathcal{L}_{L,n} &= 2\mathcal{L}_{L,n-1} - \mathcal{L}_{L,n-3}, \\ \mathcal{M}_n &= 2\mathcal{M}_{n-1} - \mathcal{M}_{n-3}. \end{aligned}$$

Let  $\psi$ ,  $\phi$  and  $\omega$  be roots of the characteristic equation  $t^3 - 2t^2 + 1 = (t^2 - t - 1)(t - 1) = 0$  corresponding to equations (2.4)–(2.6). By solving the equation, we get three distinct roots as follows:

$$(2.8) \quad \psi = \frac{1 + \sqrt{5}}{2}, \quad \phi = \frac{1 - \sqrt{5}}{2} \quad \text{and} \quad \omega = 1.$$

Note that

$$(2.9) \quad \begin{aligned} \psi + \phi + \omega &= 2 & \text{or} & \quad \psi + \phi = 1, \quad \psi - \phi = \sqrt{5} \\ \psi\phi + \psi\omega + \phi\omega &= 0 \\ \psi\phi\omega &= -1 & \text{or} & \quad \psi\phi = -1 \end{aligned}$$

and moreover  $\psi^2 + 1 = \sqrt{5}\psi$ ,  $\phi^2 + 1 = -\sqrt{5}\phi$ ,  $\psi^2 + \phi^2 = 3$ ,  $\psi^2 - \phi^2 = \sqrt{5}$  and  $\psi^3 + \phi^3 = 4$ . On the other hand,

$$(2.10) \quad \mathcal{L}_n = \frac{2(\psi^{n+1} - \phi^{n+1})}{\psi - \phi} - 1,$$

$$(2.11) \quad \mathcal{L}_{L,n} = \psi^n + \phi^n + 1,$$

$$(2.12) \quad \mathcal{M}_n = \frac{\psi^{n+2} - \phi^{n+2}}{\psi - \phi} - 1$$

are the Binet formulas of Leonardo, Leonardo–Lucas and modified Leonardo numbers, respectively. From the Binet formulas of Fibonacci and Lucas numbers, we get

$$(2.13) \quad \mathcal{L}_n = 2F_{n+1} - 1, \quad \mathcal{L}_{L,n} = L_n + 1, \quad \mathcal{M}_n = F_{n+2} - 1.$$

In addition, we have the following interrelations:

$$(2.14) \quad \begin{aligned} 5\mathcal{L}_n &= 2L_{n+1} + 4L_n - 5, & \mathcal{L}_{L,n} &= 2F_{n+1} - F_n + 1, \\ 5\mathcal{M}_n &= 3L_{n+1} + L_n - 5. \end{aligned}$$

Furthermore, there are the following relations among Leonardo, Leonardo–Lucas and modified Leonardo numbers:

$$(2.15) \quad \begin{aligned} 2\mathcal{L}_{L,n} &= \mathcal{L}_{n-2} + \mathcal{L}_n + 2, \\ 2\mathcal{M}_n &= \mathcal{L}_{n+1} - 1, \\ \mathcal{M}_n &= \mathcal{L}_{L,n} + \mathcal{L}_{L,n-1} - F_n - 3. \end{aligned}$$

In [1], the  $(-n)$ th Leonardo number with negative subscript is defined as follows:

$$(2.16) \quad \mathcal{L}_{-n} = (-1)^{-1} (\mathcal{L}_{n-2} + 1) - 1, \quad \text{for } n \geq 2.$$

Additionally, the following recurrence relation for negative subscripted Leonardo numbers is presented in [17]:

$$(2.17) \quad \mathcal{L}_{-n} = -\mathcal{L}_{-n+1} + \mathcal{L}_{-n+2} - 1, \quad \text{for } n \geq 0$$

and some values of negative subscripted Leonardo numbers are  $\mathcal{L}_{-1} = -1$ ,  $\mathcal{L}_{-2} = 1$ ,  $\mathcal{L}_{-3} = -3$ ,  $\mathcal{L}_{-4} = 3$  and  $\mathcal{L}_{-5} = -7$ .

Moreover, the matrix representation of Leonardo numbers was presented in [1] with the following  $3 \times 3$  matrix:

$$(2.18) \quad \begin{bmatrix} 2 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{bmatrix}$$

and some properties of these numbers were studied with respect to the matrix.

Now we can introduce the Fibonacci and Lucas elliptic quaternions. These were investigated by Tan et al. in [15] as follows.

The  $n$ th Fibonacci and Lucas elliptic quaternions are defined respectively by

$$(2.19) \quad \begin{aligned} \mathcal{QF}_n &= F_n + F_{n+1}e_1 + F_{n+2}e_2 + F_{n+3}e_3, \\ \mathcal{QL}_n &= L_n + L_{n+1}e_1 + L_{n+2}e_2 + L_{n+3}e_3, \end{aligned}$$

where  $F_n$  and  $L_n$  are the  $n$ th Fibonacci and Lucas numbers, respectively. Here  $e_1$ ,  $e_2$  and  $e_3$  are the basis and satisfy the relation in (1.1).

Furthermore, the Fibonacci and Lucas elliptic quaternions satisfy the following recurrence relations:

$$(2.20) \quad \begin{aligned} \mathcal{QF}_n &= \mathcal{QF}_{n-1} + \mathcal{QF}_{n-2}, \quad n \geq 2, \\ \mathcal{QL}_n &= \mathcal{QL}_{n-1} + \mathcal{QL}_{n-2}, \quad n \geq 2, \end{aligned}$$

respectively, with the initial conditions  $\mathcal{QF}_0 = e_1 + e_2 + 2e_3$ ,  $\mathcal{QF}_1 = 1 + e_1 + 2e_2 + 3e_3$  and  $\mathcal{QL}_0 = 2 + e_1 + 3e_2 + 4e_3$ ,  $\mathcal{QL}_1 = 1 + 3e_1 + 4e_2 + 7e_3$ .

**3. Leonardo, Leonardo–Lucas and modified Leonardo elliptic quaternions.** In this section, we construct Leonardo, Leonardo–Lucas and modified Leonardo elliptic quaternions and establish some properties of these number sequences. Moreover, we give matrix representations of them.

**Definition 3.1.** For  $n \geq 1$ , the  $n$ th Leonardo, Leonardo–Lucas and modified Leonardo elliptic quaternions (briefly, LEQ, LLEQ, MLEQ) are defined by

$$(3.1) \quad \begin{aligned} \mathcal{QL}_n &= \mathcal{L}_n + \mathcal{L}_{n+1}e_1 + \mathcal{L}_{n+2}e_2 + \mathcal{L}_{n+3}e_3, \\ \mathcal{QL}_{L,n} &= \mathcal{L}_{L,n} + \mathcal{L}_{L,n+1}e_1 + \mathcal{L}_{L,n+2}e_2 + \mathcal{L}_{L,n+3}e_3, \\ \mathcal{QM}_n &= \mathcal{M}_n + \mathcal{M}_{n+1}e_1 + \mathcal{M}_{n+2}e_2 + \mathcal{M}_{n+3}e_3, \end{aligned}$$

respectively, where  $e_1$ ,  $e_2$  and  $e_3$  are elliptic quaternionic units and  $\mathcal{L}_n$ ,  $\mathcal{L}_{L,n}$  and  $\mathcal{M}_n$  are the  $n$ th Leonardo, Leonardo–Lucas and modified Leonardo

numbers, respectively. The set of all LEQ, LLEQ and MLEQ are denoted by  $\mathbb{LH}$ ,  $\mathbb{LH}_L$  and  $\mathbb{MH}$ , respectively.

The sequences  $\{\mathcal{QL}_n\}$ ,  $\{\mathcal{QL}_{L,n}\}$  and  $\{\mathcal{QM}_n\}$  satisfy the following third-order linear recurrence relations, respectively:

$$(3.2) \quad \mathcal{QL}_n = 2\mathcal{QL}_{n-1} - \mathcal{QL}_{n-3},$$

$$(3.3) \quad \mathcal{QL}_{L,n} = 2\mathcal{QL}_{L,n-1} - \mathcal{QL}_{L,n-3},$$

$$(3.4) \quad \mathcal{QM}_n = 2\mathcal{QM}_{n-1} - \mathcal{QM}_{n-3}.$$

**Theorem 3.2.** For  $n \geq 2$ , LEQ, LLEQ and MLEQ satisfy the recurrence relations

$$(3.5) \quad \mathcal{QL}_n = \mathcal{QL}_{n-1} + \mathcal{QL}_{n-2} + Q^*,$$

$$(3.6) \quad \mathcal{QL}_{L,n} = \mathcal{QL}_{L,n-1} + \mathcal{QL}_{L,n-2} - Q^*,$$

$$(3.7) \quad \mathcal{QM}_n = \mathcal{QM}_{n-1} + \mathcal{QM}_{n-2} + Q^*$$

with the initial conditions

$$\begin{aligned} \mathcal{QL}_0 &= 1 + e_1 + 3e_2 + 5e_3, & \mathcal{QL}_1 &= 1 + 3e_1 + 5e_2 + 9e_3, \\ \mathcal{QL}_{L,0} &= 3 + 2e_1 + 4e_2 + 5e_3, & \mathcal{QL}_{L,1} &= 2 + 4e_1 + 5e_2 + 8e_3, \\ \mathcal{QM}_0 &= e_1 + 2e_2 + 4e_3, & \mathcal{QM}_1 &= 1 + 2e_1 + 4e_2 + 7e_3, \end{aligned}$$

where  $Q^* = 1 + e_1 + e_2 + e_3$ .

**Proof.** It can be easily proved by using the equations (2.4)–(2.6) and (3.1).  $\square$

**Theorem 3.3.** The generating functions for LEQ, LLEQ and MLEQ are

$$(3.8) \quad \begin{aligned} \text{GF}_{\mathcal{QL}_n}(t) &= \frac{\mathcal{QL}_0 + (\mathcal{QL}_1 - 2\mathcal{QL}_0)t + (\mathcal{QL}_2 - 2\mathcal{QL}_1)t^2}{t^3 - 2t + 1}, \\ \text{GF}_{\mathcal{QL}_{L,n}}(t) &= \frac{\mathcal{QL}_{L,0} + (\mathcal{QL}_{L,1} - 2\mathcal{QL}_{L,0})t + (\mathcal{QL}_{L,2} - 2\mathcal{QL}_{L,1})t^2}{t^3 - 2t + 1}, \\ \text{GF}_{\mathcal{QM}_n}(t) &= \frac{\mathcal{QM}_0 + (\mathcal{QM}_1 - 2\mathcal{QM}_0)t + (\mathcal{QM}_2 - 2\mathcal{QM}_1)t^2}{t^3 - 2t + 1}. \end{aligned}$$

**Proof.** Let  $\text{GF}_{\mathcal{QL}_n}(t)$  be the generating function for LEQ such that

$$(3.9) \quad \text{GF}_{\mathcal{QL}_n}(t) = \mathcal{QL}_0 + \mathcal{QL}_1 t + \mathcal{QL}_2 t^2 + \cdots + \mathcal{QL}_n t^n + \cdots.$$

Multiplying both sides of (3.9) by  $-2t$  and  $t^3$ , we have

$$\begin{aligned} -2t \text{GF}_{\mathcal{QL}_n}(t) &= -2\mathcal{QL}_0 t - 2\mathcal{QL}_1 t^2 - 2\mathcal{QL}_2 t^3 - \cdots - 2\mathcal{QL}_n t^{n+1} + \cdots \\ t^3 \text{GF}_{\mathcal{QL}_n}(t) &= \mathcal{QL}_0 t^3 + \mathcal{QL}_1 t^4 + \mathcal{QL}_2 t^5 + \cdots + \mathcal{QL}_n t^{n+3} + \cdots \end{aligned}$$

We have anticipated result (3.8) by using the initial conditions of LEQ and making the necessary operations. Other equations can be proved in the same way.  $\square$

**Corollary 3.4.** *Considering (2.13) and (2.20), we obtain*

$$\begin{aligned}
 \mathcal{QL}_n &= \mathcal{L}_n + \mathcal{L}_{n+1}e_1 + \mathcal{L}_{n+2}e_2 + \mathcal{L}_{n+3}e_3 \\
 (3.10) \quad &= 2F_{n+1} - 1 + (2F_{n+2} - 1)e_1 + (2F_{n+3} - 1)e_2 + (2F_{n+4} - 1)e_3 \\
 &= 2\mathcal{QF}_{n+1} - Q^*,
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{QL}_{L,n} &= \mathcal{L}_{L,n} + \mathcal{L}_{L,n+1}e_1 + \mathcal{L}_{L,n+2}e_2 + \mathcal{L}_{L,n+3}e_3 \\
 (3.11) \quad &= L_n + 1 + (L_{n+1} + 1)e_1 + (L_{n+2} + 1)e_2 + (L_{n+3} + 1)e_3 \\
 &= \mathcal{QL}_n + Q^*
 \end{aligned}$$

and

$$\begin{aligned}
 \mathcal{QM}_n &= \mathcal{M}_n + \mathcal{M}_{n+1}e_1 + \mathcal{M}_{n+2}e_2 + \mathcal{M}_{n+3}e_3 \\
 (3.12) \quad &= F_{n+2} - 1 + (F_{n+3} - 1)e_1 + (F_{n+4} - 1)e_2 + (F_{n+5} - 1)e_3 \\
 &= \mathcal{QF}_{n+2} - Q^*
 \end{aligned}$$

If we consider (2.3) and (3.10)–(3.12), we have

$$\begin{aligned}
 \mathcal{QL}_n &= 2F_{n+1} - 1 + (2F_{n+2} - 1)e_1 + (2F_{n+3} - 1)e_2 + (2F_{n+4} - 1)e_3 \\
 &= F_n + L_n + (F_{n+1} + L_{n+1})e_1 + (F_{n+2} + L_{n+2})e_2 \\
 &\quad + (F_{n+3} + L_{n+3})e_3 - Q^* \\
 &= \mathcal{QF}_n + \mathcal{QL}_n - Q^*,
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{QL}_{L,n} &= L_n + 1 + (L_{n+1} + 1)e_1 + (L_{n+2} + 1)e_2 + (L_{n+3} + 1)e_3 \\
 &= F_{n-1} + F_{n+1} + (F_n + F_{n+2})e_1 + (F_{n+1} + F_{n+3})e_2 \\
 &\quad + (F_{n+2} + F_{n+4})e_3 + Q^* \\
 &= \mathcal{QF}_{n-1} + \mathcal{QF}_{n+1} + Q^*
 \end{aligned}$$

and

$$\begin{aligned}
 \mathcal{QM}_n &= L_n + F_{n-2} - 1 + (L_{n+1} + F_{n-1} - 1)e_1 + (L_{n+2} + F_n - 1)e_2 \\
 &\quad + (L_{n+3} + F_{n+1} - 1)e_3 \\
 &= \mathcal{QL}_n + \mathcal{QF}_{n-2} - Q^*
 \end{aligned}$$

where  $\mathcal{QF}_n$  and  $\mathcal{QL}_n$  are the  $n$ th Fibonacci and Lucas elliptic quaternions, respectively.

**Corollary 3.5.** *Considering (2.14) and (2.20), we obtain*

$$\begin{aligned} 5\mathcal{QL}_n &= 5(\mathcal{L}_n + \mathcal{L}_{n+1}e_1 + \mathcal{L}_{n+2}e_2 + \mathcal{L}_{n+3}e_3) \\ &= 2\mathbb{L}_{n+1} + 4\mathbb{L}_n - 5 + (2\mathbb{L}_{n+2} + 4\mathbb{L}_{n+1} - 5)e_1 \\ &\quad + (2\mathbb{L}_{n+3} + 4\mathbb{L}_{n+2} - 5)e_2 + (2\mathbb{L}_{n+4} + 4\mathbb{L}_{n+3} - 5)e_3 \\ &= 2\mathcal{QL}_{n+1} + 4\mathcal{QL}_n - 5Q^*, \end{aligned}$$

$$\begin{aligned} \mathcal{QL}_n &= \mathcal{L}_{L,n} + \mathcal{L}_{L,n+1}e_1 + \mathcal{L}_{L,n+2}e_2 + \mathcal{L}_{L,n+3}e_3 \\ &= 2F_{n+1} - F_n + 1 + (2F_{n+2} - F_{n+1} + 1)e_1 \\ &\quad + (2F_{n+3} - F_{n+2} + 1)e_2 + (2F_{n+4} - F_{n+3} + 1)e_3 \\ &= 2\mathcal{QF}_{n+1} - \mathcal{QF}_n + Q^* \end{aligned}$$

and

$$\begin{aligned} 5\mathcal{QM}_n &= 5(\mathcal{L}_{L,n} + \mathcal{L}_{L,n+1}e_1 + \mathcal{L}_{L,n+2}e_2 + \mathcal{L}_{L,n+3}e_3) \\ &= 5\mathbb{L}_{n+1} + \mathbb{L}_n - 5 + (3\mathbb{L}_{n+2} + \mathbb{L}_{n+1} - 5)e_1 \\ &\quad + (3\mathbb{L}_{n+3} + \mathbb{L}_{n+2} - 5)e_2 + (3\mathbb{L}_{n+4} + \mathbb{L}_{n+3} - 5)e_3 \\ &= 3\mathcal{QL}_{n+1} + \mathcal{QL}_n - 5Q^*, \end{aligned}$$

where  $\mathcal{QF}_n$  and  $\mathcal{QL}_n$  are the  $n$ th Fibonacci and Lucas elliptic quaternions, respectively.

**Corollary 3.6.** *By using the equation (2.16), we can also define the negative subscripted LEQ as follows:*

$$(3.13) \quad \mathcal{QL}_{-n} = \mathcal{L}_{-n} + \mathcal{L}_{-n+1}e_1 + \mathcal{L}_{-n+2}e_2 + \mathcal{L}_{-n+3}e_3, \quad \text{for } n > 0.$$

**Theorem 3.7.** *Let  $\mathcal{QL}_{-n}$  be the  $(-n)$ th LEQ, then*

$$\begin{aligned} \mathcal{QL}_{-n} &= (-1)^n [(\mathcal{L}_{n-2} + 1) + (\mathcal{L}_{n-4} + 1)e_2] \\ &\quad + (-1)^{n-1} [(\mathcal{L}_{n-3} + 1)e_1 + (\mathcal{L}_{n-5} + 1)e_3] - Q^*, \quad \text{for } n \geq 2, \\ \mathcal{QL}_{-n} &= -\mathcal{QL}_{-n+1} + \mathcal{QL}_{-n+2} - Q^*, \quad \text{for } n > 0. \end{aligned}$$

**Proof.** It can be easily proved by using the equations (2.16) and (2.17).  $\square$

**Theorem 3.8.** *The Binet's formulas of LEQ, LLEQ and MLEQ are*

$$(3.14) \quad \mathcal{QL}_n = \frac{2\psi^{n+1}\psi^* - 2\phi^{n+1}\phi^*}{\psi - \phi} - Q^*,$$

$$(3.15) \quad \mathcal{QL}_{L,n} = \psi^n\psi^* + \phi^n\phi^* + Q^*,$$

$$(3.16) \quad \mathcal{QM}_n = \frac{\psi^{n+2}\psi^* - \phi^{n+2}\phi^*}{\psi - \phi} - Q^*$$

where  $\psi^* = 1 + \psi e_1 + \psi^2 e_2 + \psi^3 e_3$  and  $\phi^* = 1 + \phi e_1 + \phi^2 e_2 + \phi^3 e_3$ .

**Proof.** By (2.10)–(2.12) and (3.1), we obtain

$$\begin{aligned}
 Q\mathcal{L}_n &= \mathcal{L}_n + \mathcal{L}_{n+1}e_1 + \mathcal{L}_{n+2}e_2 + \mathcal{L}_{n+3}e_3 \\
 &= \frac{2(\psi^{n+1} - \phi^{n+1})}{\psi - \phi} - 1 + \left( \frac{2(\psi^{n+1} - \phi^{n+1})}{\psi - \phi} - 1 \right) e_1 \\
 &\quad + \left( \frac{2(\psi^{n+2} - \phi^{n+2})}{\psi - \phi} - 1 \right) e_2 + \left( \frac{2(\psi^{n+3} - \phi^{n+3})}{\psi - \phi} - 1 \right) e_3 \\
 &= \frac{2\psi^{n+1}}{\psi - \phi} (1 + \psi e_1 + \psi^2 e_2 + \psi^3 e_3) - \frac{2\phi^{n+1}}{\psi - \phi} (1 + \phi e_1 + \phi^2 e_2 + \phi^3 e_3) \\
 &\quad - \frac{\psi}{\psi - \phi} (1 + e_1 + e_2 + e_3) + \frac{\phi}{\psi - \phi} (1 + e_1 + e_2 + e_3) \\
 &= \frac{2\psi^{n+1}\psi^* - 2\phi^{n+1}\phi^*}{\psi - \phi} - Q^*.
 \end{aligned}$$

By (2.11) and (3.1), we get

$$\begin{aligned}
 Q\mathcal{L}_{L,n} &= \mathcal{L}_{L,n} + \mathcal{L}_{L,n+1}e_1 + \mathcal{L}_{L,n+2}e_2 + \mathcal{L}_{L,n+3}e_3 \\
 &= \psi^n + \phi^n + 1 + (\psi^{n+1} + \phi^{n+1} + 1) e_1 + (\psi^{n+2} + \phi^{n+2} + 1) e_2 \\
 &\quad + (\psi^{n+3} + \phi^{n+3} + 1) e_3 \\
 &= \psi^n (1 + \psi e_1 + \psi^2 e_2 + \psi^3 e_3) + \phi^n (1 + \phi e_1 + \phi^2 e_2 + \phi^3 e_3) + Q^* \\
 &= \psi^n \psi^* + \phi^n \phi^* + Q^*.
 \end{aligned}$$

By (2.12) and (3.1), we obtain

$$\begin{aligned}
 Q\mathcal{M}_n &= \mathcal{M}_n + \mathcal{M}_{n+1}e_1 + \mathcal{M}_{n+2}e_2 + \mathcal{M}_{n+3}e_3 \\
 &= \frac{\psi^{n+2} - \phi^{n+2}}{\psi - \phi} - 1 + \left( \frac{\psi^{n+3} - \phi^{n+3}}{\psi - \phi} - 1 \right) e_1 \\
 &\quad + \left( \frac{\psi^{n+4} - \phi^{n+4}}{\psi - \phi} - 1 \right) e_2 + \left( \frac{\psi^{n+5} - \phi^{n+5}}{\psi - \phi} - 1 \right) e_3 \\
 &= \frac{\psi^{n+2}}{\psi - \phi} (1 + \psi e_1 + \psi^2 e_2 + \psi^3 e_3) - \frac{\phi^{n+2}}{\psi - \phi} (1 + \phi e_1 + \phi^2 e_2 + \phi^3 e_3) - Q^* \\
 &= \frac{\psi^{n+2}\psi^* - \phi^{n+2}\phi^*}{\psi - \phi} - Q^*. \quad \square
 \end{aligned}$$

**Remark 3.9.** By the definitions of  $\psi^*$ ,  $\phi^*$  and  $Q^*$ , we get the following relationships:

$$\begin{aligned}
 \psi^* \phi^* &= 1 + \alpha - \beta + \gamma + \left( 1 - \sqrt{5} \frac{\Delta}{\alpha} \right) e_1 + \left( 3 - \sqrt{5} \frac{\Delta}{\beta} \right) e_2 + \left( 4 - \sqrt{5} \frac{\Delta}{\gamma} \right) e_3, \\
 \phi^* \psi^* &= 1 + \alpha - \beta + \gamma + \left( 1 + \sqrt{5} \frac{\Delta}{\alpha} \right) e_1 + \left( 3 + \sqrt{5} \frac{\Delta}{\beta} \right) e_2 + \left( 4 - \sqrt{5} \frac{\Delta}{\gamma} \right) e_3,
 \end{aligned}$$

$$\begin{aligned}\psi^*Q^* &= -1 - \psi(\alpha + \psi\beta + \psi^2\gamma) - \frac{\Delta}{\alpha}\psi e_1 + \sqrt{5}\frac{\Delta}{\beta}\psi^2 e_2 - \frac{\Delta}{\gamma}e_3 + \psi^* + Q^*, \\ \phi^*Q^* &= -1 - \phi(\alpha + \phi\beta + \phi^2\gamma) - \frac{\Delta}{\alpha}\phi e_1 + \sqrt{5}\frac{\Delta}{\beta}\phi^2 e_2 - \frac{\Delta}{\gamma}e_3 + \phi^* + Q^*, \\ Q^*\psi^* &= Q^* + \phi^* - (1 + \psi\alpha + \psi^2\beta + \psi^3\gamma) + \psi\frac{\Delta}{\alpha}e_1 + (\psi + 2)\frac{\Delta}{\beta}e_2 + \frac{\Delta}{\gamma}e_3, \\ Q^*\phi^* &= Q^* + \phi^* - (1 + \phi\alpha + \phi^2\beta + \phi^3\gamma) + \phi\frac{\Delta}{\alpha}e_1 - \frac{\Delta}{\beta}e_2 + \frac{\Delta}{\gamma}e_3.\end{aligned}$$

**Theorem 3.10.** *The exponential generating functions for LEQ, LLEQ and MLEQ are*

$$\begin{aligned}\sum_{n=0}^{\infty} \mathcal{QL}_n \frac{y^n}{n!} &= 2 \left( \frac{\psi\psi^* e^{\psi y} - \phi\phi^* e^{\phi y}}{\psi - \phi} \right) - Q^* e^y, \\ \sum_{n=0}^{\infty} \mathcal{QL}_{L,n} \frac{y^n}{n!} &= \psi^* e^{\psi y} + \phi^* e^{\phi y} + Q^* e^y, \\ \sum_{n=0}^{\infty} \mathcal{QM}_n \frac{y^n}{n!} &= \frac{\psi^{n+1}\psi^* e^{\psi y} - \phi^{n+1}\phi^* e^{\phi y}}{\psi - \phi} - Q^* e^y.\end{aligned}$$

**Proof.** By formula (3.14), we get

$$\begin{aligned}\sum_{n=0}^{\infty} \mathcal{QL}_n \frac{y^n}{n!} &= \sum_{n=0}^{\infty} \left[ 2 \left( \frac{\psi^{n+1}\psi^* - \phi^{n+1}\phi^*}{\psi - \phi} \right) - Q^* \right] \frac{y^n}{n!} \\ &= \frac{2\psi\psi^*}{\psi - \phi} \sum_{n=0}^{\infty} \frac{(\psi y)^n}{n!} - \frac{2\phi\phi^*}{\psi - \phi} \sum_{n=0}^{\infty} \frac{(\phi y)^n}{n!} - Q^* \sum_{n=0}^{\infty} \frac{y^n}{n!} \\ &= 2 \left( \frac{\psi\psi^* e^{\psi y} - \phi\phi^* e^{\phi y}}{\psi - \phi} \right) - Q^* e^y.\end{aligned}$$

Other equations can be proved in the same way. □

**Theorem 3.11.** *The Vajda's identity for the LEQ is*

$$\begin{aligned}\mathcal{QL}_{n+r} \mathcal{QL}_{n+s} - \mathcal{QL}_n \mathcal{QL}_{n+r+s} &= \frac{2\sqrt{5}}{5} (2(-1)^{n+1} \text{Fr}(\psi^s \phi^* \psi^* - \phi^s \psi^* \phi^*) \\ &\quad - (\psi^n \psi^* - \phi^n \phi^*) Q^* + Q^* (\psi^{n+s} \psi^* - \phi^{n+s} \phi^*)).\end{aligned}$$

**Proof.** Let  $\mathcal{VL}$  be the Vajda’s identity for LEQ. By the Binet formula for LEQ and the Binet formula for Fibonacci numbers, we get

$$\begin{aligned}
 \mathcal{VL} &= \mathcal{QL}_{n+r}\mathcal{QL}_{n+s} - \mathcal{QL}_n\mathcal{QL}_{n+r+s} \\
 &= \left(2\frac{\psi^{n+r+1}\psi^* - \phi^{n+r+1}\phi^*}{\psi - \phi} - Q^*\right) \left(2\frac{\psi^{n+s+1}\psi^* - \phi^{n+s+1}\phi^*}{\psi - \phi} - Q^*\right) \\
 &\quad - \left(2\frac{\psi^{n+1}\psi^* - \phi^{n+1}\phi^*}{\psi - \phi} - Q^*\right) \left(2\frac{\psi^{n+r+s+1}\psi^* - \phi^{n+r+s+1}\phi^*}{\psi - \phi} - Q^*\right) \\
 &= \frac{4}{(\psi - \phi)^2} (\psi^{2n+r+s+2} (\psi^*)^2 - \psi^{n+r+1}\phi^{n+s+1}\psi^*\phi^* \\
 &\quad - \psi^{n+s+1}\phi^{n+r+1}\phi^*\psi^* + \phi^{2n+r+s+2} (\phi^*)^2) \\
 &\quad - \frac{2}{\psi - \phi} (\psi^{n+r+1}\psi^* - \phi^{n+r+1}\phi^* + \psi^{n+1}\psi^* - \phi^{n+1}\phi^*) Q^* \\
 &\quad + \frac{2Q^*}{\psi - \phi} (\psi^{n+r+s+1}\psi^* - \phi^{n+r+s+1}\phi^* - \psi^{n+s+1}\psi^* + \phi^{n+s+1}\phi^*) \\
 &= \frac{4}{(\psi - \phi)^2} ((\psi^{n+1}\phi^{n+r+s+1} - \psi^{n+r+1}\phi^{n+s+1}) \psi^*\phi^* \\
 &\quad + (\psi^{n+r+s+1}\phi^{n+1} - \psi^{n+s+1}\phi^{n+r+1}) \phi^*\psi^*) \\
 &\quad + \frac{2}{\psi - \phi} (\psi^{n+1} (1 - \psi) \psi^* - \phi^{n+1} (1 - \phi) \phi^*) Q^* \\
 &\quad + \frac{2Q^*}{\psi - \phi} (\psi^{n+s+1} (\psi - 1) \psi^* - \phi^{n+s+1} (\phi - 1) \phi^*) \\
 &= \frac{4}{5} (-1)^{n+1} (\phi^n (\phi^r - \psi^r) \psi^*\phi^* + \psi^n (\psi^r - \phi^r) \phi^*\psi^*) \\
 &\quad + \frac{2}{\sqrt{5}} (\psi\phi (\psi^n\psi^* - \phi^n\phi^*) Q^* + Q^*\psi\phi (\phi^{n+s}\phi^* - \alpha^{n+s}\psi^*)) \\
 &= \frac{2\sqrt{5}}{5} (2(-1)^{n+1} F_r (\psi^s\phi^*\psi^* - \phi^s\psi^*\phi^*) \\
 &\quad - (\psi^n\psi^* - \phi^n\phi^*) Q^* + Q^* (\psi^{n+s}\psi^* - \phi^{n+s}\phi^*)). \quad \square
 \end{aligned}$$

In the following corollaries we have particular cases of Vajda’s identity.

**Corollary 3.12.** For  $r = -s$ , we reduce it to Catalan’s identity for LEQ as follows:

$$\begin{aligned}
 \mathcal{QL}_{n-s}\mathcal{QL}_{n+s} - \mathcal{QL}_n^2 &= \frac{2\sqrt{5}}{5} (2(-1)^{n+1} F_r (\psi^s\phi^*\psi^* - \phi^s\psi^*\phi^*) \\
 &\quad - (\psi^n\psi^* - \phi^n\phi^*) Q^* + Q^* (\psi^{n+s}\psi^* - \phi^{n+s}\phi^*)).
 \end{aligned}$$

**Corollary 3.13.** For  $s = -r = 1$ , we reduce it to Cassini identity for LEQ as follows:

$$\begin{aligned} \mathcal{QL}_{n-1}\mathcal{QL}_{n+1} - \mathcal{QL}_n^2 &= \frac{2\sqrt{5}}{5} (2(-1)^{n+1} (\psi\phi^*\psi^* - \phi\psi^*\phi^*) \\ &\quad - (\psi^n\psi^* - \phi^n\phi^*)Q^* + Q^*(\psi^{n+1}\psi^* - \phi^{n+1}\phi^*)). \end{aligned}$$

**Corollary 3.14.** For  $s \rightarrow m - n$ ,  $m \geq n$  and  $r = 1$ , we reduce it to d'Ocagne's identity for LEQ as follows:

$$\begin{aligned} \mathcal{QL}_{n+1}\mathcal{QL}_m - \mathcal{QL}_n\mathcal{QL}_{m+1} &= \frac{2\sqrt{5}}{5} (2(\psi^{m-n}\phi^*\psi^* - \phi^{m-n}\psi^*\phi^*) \\ &\quad - (\psi^n\psi^* - \phi^n\phi^*)Q^* + Q^*(\psi^m\psi^* - \phi^m\phi^*)). \end{aligned}$$

**Theorem 3.15.** The Vajda's identity for the LLEQ is

$$\begin{aligned} \mathcal{QL}_{L,n+r}\mathcal{QL}_{L,n+s} - \mathcal{QL}_{L,n}\mathcal{QL}_{L,n+r+s} \\ &= \sqrt{5}(-1)^n \phi^s F_r (\phi^s \psi^* \phi^* - \psi^s \phi^* \psi^*) + \psi^n (\psi^r - 1) (\psi^* Q^* - \psi^s Q^* \psi^*) \\ &\quad + \phi^n (\phi^r - 1) (\phi^* Q^* - \phi^s Q^* \phi^*). \end{aligned}$$

**Proof.** Let  $\mathcal{VL}_L$  be the Vajda's identity for LLEQ. By the Binet formula for LLEQ and the Binet formula for Fibonacci numbers, we get

$$\begin{aligned} \mathcal{VL}_L &= \mathcal{QL}_{L,n+r}\mathcal{QL}_{L,n+s} - \mathcal{QL}_{L,n}\mathcal{QL}_{L,n+r+s} \\ &= (\psi^{n+r}\psi^* + \phi^{n+r}\phi^* + Q^*) (\psi^{n+s}\psi^* + \phi^{n+s}\phi^* + Q^*) \\ &\quad - (\psi^n\psi^* + \phi^n\phi^* + Q^*) (\psi^{n+r+s}\psi^* + \phi^{n+r+s}\phi^* + Q^*) \\ &= (\psi^{n+r}\phi^{n+s} - \psi^n\phi^{n+r+s}) \psi^* \phi^* + (\psi^{n+s}\phi^{n+r} - \psi^{n+r+s}\phi^n) \phi^* \psi^* \\ &\quad + (\psi^{n+r} - \psi^n) \psi^* Q^* + (\psi^{n+s} - \psi^{n+r+s}) Q^* \psi^* \\ &\quad + (\phi^{n+r} - \phi^n) \phi^* Q^* + (\phi^{n+s} - \phi^{n+r+s}) Q^* \phi^* \\ &= \psi^n \phi^{n+s} (\psi^r - \phi^r) \psi^* \phi^* + \psi^{n+s} \phi^n (\phi^r - \psi^r) \phi^* \psi^* + \psi^n (\psi^r - 1) \psi^* Q^* \\ &\quad + \psi^{n+s} (1 - \psi^r) Q^* \psi^* + \phi^n (\phi^r - 1) \phi^* Q^* + \phi^{n+s} (1 - \phi^r) Q^* \phi^* \\ &= \sqrt{5}(-1)^n \phi^s F_r \psi^* \phi^* - \sqrt{5}(-1)^n \psi^s F_r \phi^* \psi^* + \psi^n (\psi^r - 1) \psi^* Q^* \\ &\quad + \psi^{n+s} (1 - \psi^r) Q^* \psi^* + \phi^n (\phi^r - 1) \phi^* Q^* + \phi^{n+s} (1 - \phi^r) Q^* \phi^* \\ &= \sqrt{5}(-1)^n \phi^s F_r (\phi^s \psi^* \phi^* - \psi^s \phi^* \psi^*) + \psi^n (\psi^r - 1) (\psi^* Q^* - \psi^s Q^* \psi^*) \\ &\quad + \phi^n (\phi^r - 1) (\phi^* Q^* - \phi^s Q^* \phi^*). \quad \square \end{aligned}$$

In the following corollaries we have particular cases of Vajda’s identity.

**Corollary 3.16.** *For  $r = -s$ , we reduce it to Catalan’s identity for LLEQ as follows:*

$$\begin{aligned} & \mathcal{QL}_{L,n-s}\mathcal{QL}_{L,n+s} - \mathcal{QL}_{L,n}^2 \\ &= \sqrt{5}(-1)^n \phi^s (\phi^s \psi^* \phi^* - \psi^s \phi^* \psi^*) + \psi^n (\psi^{-s} - 1) (\psi^* Q^* - \psi^s Q^* \psi^*) \\ & \quad + \phi^n (\phi^{-s} - 1) (\phi^* Q^* - \phi^s Q^* \phi^*). \end{aligned}$$

**Corollary 3.17.** *For  $s = -r = 1$ , we reduce it to Cassini identity for LLEQ as follows:*

$$\begin{aligned} & \mathcal{QL}_{L,n-1}\mathcal{QL}_{L,n+1} - \mathcal{QL}_{L,n}^2 \\ &= \sqrt{5}(-1)^n \phi (\phi \psi^* \phi^* - \psi \phi^* \psi^*) - \left(\frac{3 - \sqrt{5}}{2}\right) \psi^n (\psi^* Q^* - \psi Q^* \psi^*) \\ & \quad - \phi \left(\frac{3 + \sqrt{5}}{2}\right) (\phi^* Q^* - \phi Q^* \phi^*). \end{aligned}$$

**Corollary 3.18.** *For  $s \rightarrow m - n$ ,  $m \geq n$  and  $r = 1$ , we reduce it to d’Ocagne’s identity for LLEQ as follows:*

$$\begin{aligned} & \mathcal{QL}_{L,n+1}\mathcal{QL}_{L,m} - \mathcal{QL}_{L,n}\mathcal{QL}_{L,m+1} \\ &= \sqrt{5}(-1)^n \phi^{m-n} (\phi^{m-n} \psi^* \phi^* - \psi^{m-n} \phi^* \psi^*) + \psi^n \phi (\psi^* Q^* - \psi^{m-n} Q^* \psi^*) \\ & \quad - \phi^n \psi (\phi^* Q^* - \phi^{m-n} Q^* \phi^*). \end{aligned}$$

**Theorem 3.19.** *The Vajda’s identity for the MLEQ is*

$$\begin{aligned} & \mathcal{QM}_{n+r}\mathcal{QM}_{n+s} - \mathcal{QM}_n\mathcal{QM}_{n+r+s} \\ &= (-1)^n F_r (\psi^s \phi^* \psi^* - \phi^s \psi^* \phi^*) + \frac{1}{\sqrt{5}} \psi^{n+2} (1 - \psi^r) (\psi^* Q^* - \psi^s Q^* \psi^*) \\ & \quad - \frac{1}{\sqrt{5}} \phi^{n+2} (1 - \phi^r) (\phi^* Q^* - \phi^s Q^* \phi^*). \end{aligned}$$

**Proof.** Let  $\mathcal{VM}$  be the Vajda's identity for MLEQ. By the Binet formula for MLEQ and the Binet formula for Fibonacci numbers, we get

$$\begin{aligned}
\mathcal{VM} &= \mathcal{QM}_{n+r} \mathcal{QM}_{n+s} - \mathcal{QM}_n \mathcal{QM}_{n+r+s} \\
&= \left( \frac{\psi^{n+r+2} \psi^* - \phi^{n+r+2} \phi^*}{\psi - \phi} - Q^* \right) \left( \frac{\psi^{n+s+2} \psi^* - \phi^{n+s+2} \phi^*}{\psi - \phi} - Q^* \right) \\
&\quad - \left( \frac{\psi^{n+2} \psi^* - \phi^{n+2} \phi^*}{\psi - \phi} - Q^* \right) \left( \frac{\psi^{n+r+s+2} \psi^* - \phi^{n+r+s+2} \phi^*}{\psi - \phi} - Q^* \right) \\
&= \frac{1}{(\psi - \phi)^2} \left( (\psi^{n+2} \phi^{n+r+s+2} - \psi^{n+r+2} \phi^{n+s+2}) \psi^* \phi^* \right. \\
&\quad \left. + (\psi^{n+r+s+2} \phi^{n+2} - \psi^{n+s+2} \phi^{n+r+2}) \phi^* \psi^* \right) \\
&\quad + \frac{1}{\psi - \phi} \left( \psi^{n+2} (1 - \psi^r) \psi^* + \phi^{n+2} (\phi^r - 1) \phi^* \right) Q^* \\
&\quad + \frac{Q^*}{\psi - \phi} \left( \psi^{n+s+2} (\psi^r - 1) \psi^* + \phi^{n+s+2} (1 - \phi^r) \phi^* \right) \\
&= \frac{\psi^r \phi^r}{(\psi - \phi)^2} \psi^{n+2} \phi^{n+2} (\psi^s \phi^* \psi^* - \phi^s \psi^* \phi^*) \\
&\quad + \frac{1 - \psi^r}{\psi - \phi} (\psi^{n+2} \psi^* Q^* - \psi^{n+s+2} Q^* \psi^*) \\
&\quad + \frac{1 - \phi^r}{\psi - \phi} (\phi^{n+2} Q^* \phi^* - \psi^{n+2} \phi^* Q^*) \\
&= (-1)^n F_r (\psi^s \phi^* \psi^* - \phi^s \psi^* \phi^*) + \frac{1}{\sqrt{5}} \psi^{n+2} (1 - \psi^r) (\psi^* Q^* - \psi^s Q^* \psi^*) \\
&\quad - \frac{1}{\sqrt{5}} \phi^{n+2} (1 - \phi^r) (\phi^* Q^* - \phi^s Q^* \phi^*). \quad \square
\end{aligned}$$

In the following corollaries we have particular cases of Vajda's identity.

**Corollary 3.20.** *For  $r = -s$ , we reduce it to Catalan's identity for MLEQ as follows:*

$$\begin{aligned}
\mathcal{QM}_{n-s} \mathcal{QM}_{n+s} - \mathcal{QM}_n^2 &= (-1)^{n+s+3} F_s (\psi^s \phi^* \psi^* - \phi^s \psi^* \phi^*) \\
&\quad + \frac{1}{\sqrt{5}} \psi^{n+2} (1 - \psi^{-s}) (\psi^* Q^* - \psi^s Q^* \psi^*) \\
&\quad - \frac{1}{\sqrt{5}} \phi^{n+2} (1 - \phi^{-s}) (\phi^* Q^* - \phi^s Q^* \phi^*).
\end{aligned}$$

**Corollary 3.21.** *For  $s = -r = 1$ , we reduce it to Cassini identity for MLEQ as follows:*

$$\begin{aligned} \mathcal{QM}_{n-1}\mathcal{QM}_{n+1} - \mathcal{QM}_n^2 &= (-1)^n (\psi\phi^*\psi^* - \phi\psi^*\phi^*) \\ &\quad + \frac{1}{\sqrt{5}}\psi^{n+2} (1 - \psi^{-1}) (\psi^*Q^* - \psi Q^*\psi^*) \\ &\quad - \frac{1}{\sqrt{5}}\phi^{n+2} (1 - \phi^{-1}) (\phi^*Q^* - \phi Q^*\phi^*). \end{aligned}$$

**Corollary 3.22.** *For  $s \rightarrow m - n$ ,  $m \geq n$  and  $r = 1$ , we it reduce to d’Ocagne’s identity for MLEQ as follows:*

$$\begin{aligned} \mathcal{QM}_{n+1}\mathcal{QM}_m - \mathcal{QM}_n\mathcal{QM}_{m+1} &= (-1)^n (\psi^{m-n}\phi^*\psi^* - \phi^{m-n}\psi^*\phi^*) \\ &\quad + \frac{1}{\sqrt{5}}\psi^{n+2}\phi (\psi^*Q^* - \psi^{m-n}Q^*\psi^*) \\ &\quad - \frac{1}{\sqrt{5}}\phi^{n+2}\psi (\phi^*Q^* - \phi^{m-n}Q^*\phi^*). \end{aligned}$$

**Corollary 3.23.** *Considering (2.13) and (2.14), we obtain*

$$\begin{aligned} 2\mathcal{QM}_n &= 2(\mathcal{M}_n + \mathcal{M}_{n+1}e_1 + \mathcal{M}_{n+2}e_2 + \mathcal{M}_{n+3}e_3) \\ &= 2\mathcal{L}_{n+1} - 1 + (2\mathcal{L}_{n+2} - 1)e_1 + (2\mathcal{L}_{n+3} - 1)e_2 + (2\mathcal{L}_{n+4} - 1)e_3 \\ &= 2\mathcal{QL}_{n+1} - Q^*, \end{aligned}$$

$$\begin{aligned} 2\mathcal{QL}_{L,n} &= 2(\mathcal{L}_{L,n} + \mathcal{L}_{L,n+1}e_1 + \mathcal{L}_{L,n+2}e_2 + \mathcal{L}_{L,n+3}e_3) \\ &= \mathcal{L}_{L,n-2} + \mathcal{L}_{L,n} + 2 + (\mathcal{L}_{L,n-1} + \mathcal{L}_{L,n+1} + 2)e_1 \\ &\quad + (\mathcal{L}_{L,n} + \mathcal{L}_{L,n+2} + 2)e_2 + (\mathcal{L}_{L,n+1} + \mathcal{L}_{L,n+3} + 2)e_3 \\ &= 2\mathcal{QL}_{n-2} + \mathcal{QL}_n + 2Q^* \end{aligned}$$

and

$$\begin{aligned} \mathcal{QM}_n &= \mathcal{M}_n + \mathcal{M}_{n+1}e_1 + \mathcal{M}_{n+2}e_2 + \mathcal{M}_{n+3}e_3 \\ &= \mathcal{L}_{L,n} + \mathcal{L}_{L,n-1} - F_n - 3 + (\mathcal{L}_{L,n+1} + \mathcal{L}_{L,n} - F_{n+1} - 3)e_1 \\ &\quad + (\mathcal{L}_{L,n+2} + \mathcal{L}_{L,n+1} - F_{n+2} - 3)e_2 \\ &\quad + (\mathcal{L}_{L,n+3} + \mathcal{L}_{L,n+2} - F_{n+3} - 3)e_3 \\ &= \mathcal{QL}_{L,n} + \mathcal{QL}_{L,n-1} - \mathcal{QF}_n - 3Q^*, \end{aligned}$$

where  $\mathcal{QF}_n$  is the  $n$ th Fibonacci elliptic quaternion.

**4. The matrix representations of LEQ, LLEQ and MLEQ.** In addition to previous formulations, we can define the matrix equalities for LEQ, LLEQ, and MLEQ using the following expressions.

**Theorem 4.1.** *Let  $\mathcal{QL}_n$ ,  $\mathcal{QL}_{L,n}$  and  $\mathcal{QM}_n$  be the  $n$ th LEQ, LLEQ and MLEQ, respectively. For every  $n > 0$ , the following matrix equalities hold:*

$$\begin{bmatrix} \mathcal{QL}_3 & \mathcal{QL}_2 & \mathcal{QL}_1 \\ \mathcal{QL}_2 & \mathcal{QL}_1 & \mathcal{QL}_0 \\ \mathcal{QL}_1 & \mathcal{QL}_0 & \mathcal{QL}_{-1} \end{bmatrix} \begin{bmatrix} 2 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{bmatrix}^n = \begin{bmatrix} \mathcal{QL}_{n+3} & \mathcal{QL}_{n+2} & \mathcal{QL}_{n+1} \\ \mathcal{QL}_{n+2} & \mathcal{QL}_{n+1} & \mathcal{QL}_n \\ \mathcal{QL}_{n+1} & \mathcal{QL}_n & \mathcal{QL}_{n-1} \end{bmatrix},$$

$$\begin{bmatrix} \mathcal{QL}_{L,3} & \mathcal{QL}_{L,2} & \mathcal{QL}_{L,1} \\ \mathcal{QL}_{L,2} & \mathcal{QL}_{L,1} & \mathcal{QL}_{L,0} \\ \mathcal{QL}_{L,1} & \mathcal{QL}_{L,0} & \mathcal{QL}_{L,-1} \end{bmatrix} \begin{bmatrix} 2 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{bmatrix}^n = \begin{bmatrix} \mathcal{QL}_{L,n+3} & \mathcal{QL}_{L,n+2} & \mathcal{QL}_{L,n+1} \\ \mathcal{QL}_{L,n+2} & \mathcal{QL}_{L,n+1} & \mathcal{QL}_{L,n} \\ \mathcal{QL}_{L,n+1} & \mathcal{QL}_{L,n} & \mathcal{QL}_{L,n-1} \end{bmatrix},$$

$$\begin{bmatrix} \mathcal{QM}_3 & \mathcal{QM}_2 & \mathcal{QM}_1 \\ \mathcal{QM}_2 & \mathcal{QM}_1 & \mathcal{QM}_0 \\ \mathcal{QM}_1 & \mathcal{QM}_0 & \mathcal{QM}_{-1} \end{bmatrix} \begin{bmatrix} 2 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{bmatrix}^n = \begin{bmatrix} \mathcal{QM}_{n+3} & \mathcal{QM}_{n+2} & \mathcal{QM}_{n+1} \\ \mathcal{QM}_{n+2} & \mathcal{QM}_{n+1} & \mathcal{QM}_n \\ \mathcal{QM}_{n+1} & \mathcal{QM}_n & \mathcal{QM}_{n-1} \end{bmatrix}$$

and

$$\begin{bmatrix} \mathcal{QL}_{n+2} \\ \mathcal{QL}_{n+1} \\ \mathcal{QL}_n \end{bmatrix} = \begin{bmatrix} 2 & 0 & -1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}^n \begin{bmatrix} \mathcal{QL}_2 \\ \mathcal{QL}_1 \\ \mathcal{QL}_0 \end{bmatrix},$$

$$\begin{bmatrix} \mathcal{QL}_{L,n+2} \\ \mathcal{QL}_{L,n+1} \\ \mathcal{QL}_{L,n} \end{bmatrix} = \begin{bmatrix} 2 & 0 & -1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}^n \begin{bmatrix} \mathcal{QL}_{L,2} \\ \mathcal{QL}_{L,1} \\ \mathcal{QL}_{L,0} \end{bmatrix},$$

$$\begin{bmatrix} \mathcal{QM}_{n+2} \\ \mathcal{QM}_{n+1} \\ \mathcal{QM}_n \end{bmatrix} = \begin{bmatrix} 2 & 0 & -1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}^n \begin{bmatrix} \mathcal{QM}_2 \\ \mathcal{QM}_1 \\ \mathcal{QM}_0 \end{bmatrix}.$$

**Proof.** The proof is obvious by mathematical induction on  $n$ . The equality hold for  $n = 1$ . Now suppose that the equality is true for  $n > 1$ . Then we can verify it for  $n + 1$  as follows

$$\begin{aligned} & \begin{bmatrix} \mathcal{QL}_3 & \mathcal{QL}_2 & \mathcal{QL}_1 \\ \mathcal{QL}_2 & \mathcal{QL}_1 & \mathcal{QL}_0 \\ \mathcal{QL}_1 & \mathcal{QL}_0 & \mathcal{QL}_{-1} \end{bmatrix} \begin{bmatrix} 2 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{bmatrix}^{n+1} \\ &= \begin{bmatrix} \mathcal{QL}_3 & \mathcal{QL}_2 & \mathcal{QL}_1 \\ \mathcal{QL}_2 & \mathcal{QL}_1 & \mathcal{QL}_0 \\ \mathcal{QL}_1 & \mathcal{QL}_0 & \mathcal{QL}_{-1} \end{bmatrix} \begin{bmatrix} 2 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{bmatrix}^n \begin{bmatrix} 2 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} \mathcal{QL}_{n+3} & \mathcal{QL}_{n+2} & \mathcal{QL}_{n+1} \\ \mathcal{QL}_{n+2} & \mathcal{QL}_{n+1} & \mathcal{QL}_n \\ \mathcal{QL}_{n+1} & \mathcal{QL}_n & \mathcal{QL}_{n-1} \end{bmatrix} \begin{bmatrix} 2 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} \mathcal{QL}_{n+4} & \mathcal{QL}_{n+3} & \mathcal{QL}_{n+2} \\ \mathcal{QL}_{n+3} & \mathcal{QL}_{n+2} & \mathcal{QL}_{n+1} \\ \mathcal{QL}_{n+2} & \mathcal{QL}_{n+1} & \mathcal{QL}_n \end{bmatrix}. \quad \square \end{aligned}$$

**Corollary 4.2.** Consider  $\mathcal{QL}_n, \mathcal{QL}_m \in \mathbb{LH}$ . We have the following:

$$\begin{aligned} \varphi_{\mathcal{QL}_n} : \mathbb{LH} &\longrightarrow \mathbb{LH} \\ \mathcal{QL}_m &\longmapsto \varphi_{\mathcal{QL}_n}(\mathcal{QL}_m) = \mathcal{QL}_n \mathcal{QL}_m \end{aligned}$$

and

$$\begin{aligned} \mu_{\mathcal{QL}_n} : \mathbb{LH} &\longrightarrow \mathbb{LH} \\ \mathcal{QL}_m &\longmapsto \mu_{\mathcal{QL}_n}(\mathcal{QL}_m) = \mathcal{QL}_m \mathcal{QL}_n, \end{aligned}$$

where  $\varphi_{\mathcal{QL}_n}$  and  $\mu_{\mathcal{QL}_n}$  are respectively given as

$$\begin{aligned} \mathcal{L}\mathcal{QL}_n &= \begin{bmatrix} \mathcal{L}_n & -\alpha\mathcal{L}_{n+1} & -\beta\mathcal{L}_{n+2} & -\gamma\mathcal{L}_{n+3} \\ \mathcal{L}_{n+1} & \mathcal{L}_n & -\frac{\Delta}{\alpha}\mathcal{L}_{n+3} & \frac{\Delta}{\alpha}\mathcal{L}_{n+2} \\ \mathcal{L}_{n+2} & \frac{\Delta}{\beta}\mathcal{L}_{n+3} & \mathcal{L}_n & -\frac{\Delta}{\beta}\mathcal{L}_{n+1} \\ \mathcal{L}_{n+3} & -\frac{\Delta}{\gamma}\mathcal{L}_{n+2} & \frac{\Delta}{\gamma}\mathcal{L}_{n+1} & \mathcal{L}_n \end{bmatrix}, \\ \mathcal{R}\mathcal{QL}_n &= \begin{bmatrix} \mathcal{L}_n & -\alpha\mathcal{L}_{n+1} & -\beta\mathcal{L}_{n+2} & -\gamma\mathcal{L}_{n+3} \\ \mathcal{L}_{n+1} & \mathcal{L}_n & \frac{\Delta}{\alpha}\mathcal{L}_{n+3} & -\frac{\Delta}{\alpha}\mathcal{L}_{n+2} \\ \mathcal{L}_{n+2} & -\frac{\Delta}{\beta}\mathcal{L}_{n+3} & \mathcal{L}_n & \frac{\Delta}{\beta}\mathcal{L}_{n+1} \\ \mathcal{L}_{n+3} & \frac{\Delta}{\gamma}\mathcal{L}_{n+2} & -\frac{\Delta}{\gamma}\mathcal{L}_{n+1} & \mathcal{L}_n \end{bmatrix}. \end{aligned}$$

Other representations for the numbers in  $\mathbb{LHL}$  and  $\mathbb{MLH}$  can be shown in the same way.

**5. Conclusions.** Elliptic quaternions defined in [10] has been given for a special case by replacing real number coefficients by Leonardo, Leonardo–Lucas and modified Leonardo number coefficients. We first defined elementary operations and its  $4 \times 4$  matrix representations on the set of elliptic quaternions. Then we gave the definitions and relationships of Leonardo, Leonardo–Lucas and modified Leonardo numbers. We then went on to investigate relationships, recurrence relation and Binet formula of Leonardo, Leonardo–Lucas and modified Leonardo elliptic quaternions. We also provided the generating function. We finally investigated Vajda’s identity and gave some corollaries for special cases such as Catalan, Cassini and D’Ocagne’s identities. Lastly, we studied two different matrix representations of these numbers.

In the following table (Table 2), we show the numbers which are mentioned in this paper.

Number	Definition
Elliptic Quaternions ( $\mathbb{H}_{\alpha,\beta,\gamma}$ ) [10]	$Q = q_0e_0 + q_1e_1 + q_2e_2 + q_3e_3 :$ $q_0, q_1, q_2, q_3 \in \mathbb{R}$
$n$ th Fibonacci Numbers	$F_n = F_{n-1} + F_{n-2}, F_0 = 0, F_1 = 1$
$n$ th Lucas Numbers	$L_n = L_{n-1} + L_{n-2}, L_0 = 2, L_1 = 1$
$n$ th Leonardo Numbers	$\mathcal{L}_n = \mathcal{L}_{n-1} + \mathcal{L}_{n-2} + 1,$ $n \geq 2, \mathcal{L}_0 = \mathcal{L}_1 = 1$
$n$ th Leonardo-Lucas Numbers	$\mathcal{L}_{L,n} = \mathcal{L}_{L,n-1} + \mathcal{L}_{L,n-2} + 1,$ $n \geq 2, \mathcal{L}_{L,0} = 3, \mathcal{L}_{L,1} = 2$
$n$ th modified Leonardo Numbers	$\mathcal{M}_n = \mathcal{M}_{n-1} + \mathcal{M}_{n-2} + 1,$ $n \geq 2, \mathcal{M}_0 = 0, \mathcal{M}_1 = 1$
$n$ th Fibonacci Elliptic Quaternions	$\mathcal{Q}F_n = F_n + F_{n+1}e_1 + F_{n+2}e_2 + F_{n+3}e_3$
$n$ th Lucas Elliptic Quaternions	$\mathcal{Q}L_n = L_n + L_{n+1}e_1 + L_{n+2}e_2 + L_{n+3}e_3$
$n$ th Leonardo Elliptic Quaternions ( $\mathbb{L}\mathbb{H}$ )	$\mathcal{Q}\mathcal{L}_n = 2\mathcal{Q}\mathcal{L}_{n-1} - \mathcal{Q}\mathcal{L}_{n-3}$
$n$ th Leonardo-Lucas Elliptic Quaternions ( $\mathbb{L}\mathbb{H}_L$ )	$\mathcal{Q}\mathcal{L}_{L,n} = 2\mathcal{Q}\mathcal{L}_{L,n-1} - \mathcal{Q}\mathcal{L}_{L,n-3}$
$n$ th Modified-Leonardo Elliptic Quaternions ( $\mathbb{M}\mathbb{H}$ )	$\mathcal{Q}\mathcal{M}_n = 2\mathcal{Q}\mathcal{M}_{n-1} - \mathcal{Q}\mathcal{M}_{n-3}$

TABLE 2. The numbers which are mentioned in this paper.

## REFERENCES

- [1] Alp, Y., Koçer, E. G., *Some properties of Leonardo numbers*. Konuralp J. Math. **9**(1) (2021), 183–189.
- [2] Catarino, P., Borges, A., *A note on incomplete Leonardo numbers*, Integers **20** (2020), Paper No. A43, 7 pp.
- [3] Catarino, P. M., Borges, A., *On Leonardo numbers*, Acta Math. Univ. Comenian. (N.S.) **89**(1) (2019), 75–86.
- [4] Çolakoğlu, H. B., Özdemir, M., *Generalized elliptical quaternions with some applications*. Turkish J. Math. **47**(1) (2023), 351–371.
- [5] Dijkstra, E., *Archive: Fibonacci numbers and Leonardo numbers (ewd 797)*, 1981.
- [6] Isbilir, Z., Akyigit, M., Tosun, M., *Pauli-Leonardo quaternions*, Notes on Number Theory and Discrete Mathematics **29**(1) (2023), 1–16.
- [7] Karataş, A., *On complex Leonardo numbers*, Notes on Number Theory and Discrete Mathematics **28**(3) (2022), 458–465.
- [8] Kuhapatanakul, K., Ruankong, P., *On generalized Leonardo  $p$ -numbers*, J. Integer Seq. **27**(4) (2024), Art. 24.4.6, 10 pp.

- [9] Mangueira, M., Vieira, R., Alves, F., Catarino, P., *Leonardo's bivariate and complex polynomials*, Notes on Number Theory and Discrete Mathematics **28**(1) (2022), 115–123.
- [10] Özdemir, M., *An alternative approach to elliptical motion*, Adv. Appl. Clifford Algebr. **26** (2016), 279–304.
- [11] Prasad, K., Mahato, H., Kumari, M., Mohanty, R., *On the generalized Leonardo Pisano octonions*, Nat. Acad. Sci. Lett. **47**(5) (2024), 579–585.
- [12] Prasad, K., Mohanty, R., Kumari, M., Mahato, H., *Some new families of generalized  $k$ -Leonardo and Gaussian Leonardo numbers*, Commun. Comb. Optim. **9**(3) (2024), 539–553.
- [13] Rahebi, M., Yayli, Y., *Elliptic quaternion and elliptic linear interpolation*, Politeknik Dergisi **27**(3) (2024), 1189–1195.
- [14] Soykan, Y., *Generalized Leonardo numbers*, Journal of Progressive Research in Mathematics **18**(4) (2021), 58–84.
- [15] Tan, E., Yaman, T., Gök, I., *On Fibonacci and Lucas elliptic quaternions*, 2024, to appear.
- [16] Vieira, R. P. M., Alves, F. R. V., Catarino, P. M. M. C., *Relações bidimensionais identidades da sequência de Leonardo*, Revista Sergipana de Matemática e Educação Matemática **4**(2) (2019), 156–173.
- [17] Vieira, R. P. M., dos Santos Mangueira, M. C., Alves, F. R. V., Catarino, P. M. M. C., *A forma matricial dos números de Leonardo*, Ciencia e natura **42** (2020), e100–e100.

Furkan Seçgin  
Department of Mathematics  
Ankara University  
06100 Beşevler, Ankara  
Turkey  
e-mail: [fsecgin@ankara.edu.tr](mailto:fsecgin@ankara.edu.tr)

Received December 12, 2024.