

PAWEŁ BEDNARZ, ANNA KOSIOROWSKA
and ADRIAN MICHALSKI

Special bihyperbolic numbers and their connections with triangular tables and matrices

ABSTRACT. In this paper we express special bihyperbolic numbers as parade-terminants and parapermanents of some triangular matrices. Moreover, by applying the connections between these parameters of triangular tables and the determinants and permanents of lower Hessenberg matrices, we obtain another expressions of these numbers, using matrices which are not triangular.

1. Introduction. Let $n \geq 0$ be an integer. The n th balancing number B_n , Lucas-balancing number C_n , Mersenne number M_n and Mersenne–Lucas number H_n are given by the following recursive definitions:

$$B_n = 6B_{n-1} - B_{n-2}, \text{ for } n \geq 2 \text{ with } B_0 = 0, B_1 = 1,$$

$$C_n = 6C_{n-1} - C_{n-2}, \text{ for } n \geq 2 \text{ with } C_0 = 1, C_1 = 3,$$

$$M_n = 3M_{n-1} - 2M_{n-2}, \text{ for } n \geq 2 \text{ with } M_0 = 0, M_1 = 1,$$

$$H_n = 3H_{n-1} - 2H_{n-2}, \text{ for } n \geq 2 \text{ with } H_0 = 2, H_1 = 3.$$

In Table 1 we list first numbers of sequences defined above.

2010 *Mathematics Subject Classification.* 15A15, 11B39, 47B36.

Key words and phrases. Bihyperbolic numbers, balancing numbers, Mersenne numbers, triangular matrix, Hessenberg matrix.

n	0	1	2	3	4	5	5	6
B_n	0	1	6	35	204	1189	6930	40391
C_n	1	3	17	99	577	3363	19601	114243
M_n	0	1	3	7	15	31	63	127
H_n	2	3	5	9	17	33	65	129

TABLE 1. The first terms of sequences B_n, C_n, M_n, H_n .

Balancing numbers were introduced by Behera and Panda in [2]. Later, in [18], Panda defined Lucas-balancing numbers. These two sequences were extensively studied, also by considering some generalizations. For more details, see for example [8, 11, 20]. The literature on Mersenne and Mersenne–Lucas sequences is also broad, see [4, 9, 16, 22] among others.

For the integer sequences considered in this paper: balancing numbers, Lucas-balancing numbers, Mersenne numbers and Mersenne–Lucas numbers, we also provide their corresponding entries in the OEIS (The On-Line Encyclopedia of Integer Sequences):

$$B_n : \text{A001109}, \quad C_n : \text{A001541}, \quad M_n : \text{A000225}, \quad H_n : \text{A000051}.$$

The hyperbolic unit j was introduced in 1848 by J. Cockle in [10]. Elements of the set $\mathbb{H} = \{a + bj : a, b \in \mathbb{R}, j^2 = 1, j \neq \pm 1\}$ are known as *hyperbolic numbers*. The hyperbolic numbers, which are known also as *split complex numbers* or *double numbers* were studied for the first time in the 19th century. Originally, the non-Euclidean framework was described by Yaglom [23], while a systematic modern treatment of hypercomplex systems was given by Olariu [17] and by Kantor and Solodovnikov [15]. Fundamental algebraic properties of hyperbolic numbers were studied in detail by Rochon and Shapiro [21].

One of the generalizations of hyperbolic numbers was introduced in [19]. Let us denote by \mathbb{H}_2 the set of all numbers of the form

$$\zeta = x_0 + j_1x_1 + j_2x_2 + j_3x_3,$$

where $x_0, x_1, x_2, x_3 \in \mathbb{R}$ and operators $j_1, j_2, j_3 \notin \mathbb{R}$ satisfy conditions

$$j_1^2 = j_2^2 = j_3^2 = 1, \quad j_1j_2 = j_2j_1 = j_3, \quad j_1j_3 = j_3j_1 = j_2, \quad j_2j_3 = j_3j_2 = j_1.$$

The elements of the set \mathbb{H}_2 are called *bihyperbolic numbers*.

Addition and multiplication of bihyperbolic numbers are performed analogously to algebraic expressions. These operations are associative and commutative on \mathbb{H}_2 . Moreover, multiplication is distributive over addition, so $(\mathbb{H}_2, +, \cdot)$ is a commutative ring.

Bihyperbolic numbers are well known in the literature, their properties can be found for example in [3, 21]. Recently, bihyperbolic extensions of

classical integer sequences have also been investigated; see, for example, the bihyperbolic Tribonacci-type sequences introduced in [14].

Some special cases of bihyperbolic numbers were studied in the literature. In particular, in [7], authors defined bihyperbolic numbers of the Fibonacci type. Later, following this research, other types of bihyperbolic numbers were introduced and we will focus on four of them:

- bihyperbolic balancing numbers and bihyperbolic Lucas-balancing numbers defined in [6],
- bihyperbolic Mersenne numbers and bihyperbolic Mersenne–Lucas numbers defined in [5].

Let $n \geq 0$ be an integer. The n th bihyperbolic balancing number BhB_n , bihyperbolic Lucas-balancing number BhC_n , bihyperbolic Mersenne number BhM_n and bihyperbolic Mersenne–Lucas number BhH_n are defined in the following way:

$$\begin{aligned}
 (1) \quad & BhB_n = B_n + j_1 B_{n+1} + j_2 B_{n+2} + j_3 B_{n+3}, \\
 (2) \quad & BhC_n = C_n + j_1 C_{n+1} + j_2 C_{n+2} + j_3 C_{n+3}, \\
 (3) \quad & BhM_n = M_n + j_1 M_{n+1} + j_2 M_{n+2} + j_3 M_{n+3}, \\
 (4) \quad & BhH_n = H_n + j_1 H_{n+1} + j_2 H_{n+2} + j_3 H_{n+3},
 \end{aligned}$$

where B_n is the n th balancing number, C_n is the n th Lucas-balancing number, M_n is the n th Mersenne number and H_n is the n th Mersenne–Lucas number.

Below we list first four terms of each sequence mentioned above.

$$\begin{aligned}
 & BhB_0 = j_1 + 6j_2 + 35j_3, \\
 & BhB_1 = 1 + 6j_1 + 35j_2 + 204j_3, \\
 (5) \quad & BhB_2 = 6 + 35j_1 + 204j_2 + 1189j_3, \\
 & BhB_3 = 35 + 204j_1 + 1189j_2 + 6930j_3, \\
 & \quad \quad \quad \vdots \\
 & BhC_0 = 1 + 3j_1 + 17j_2 + 99j_3, \\
 & BhC_1 = 3 + 17j_1 + 99j_2 + 577j_3, \\
 (6) \quad & BhC_2 = 17 + 99j_1 + 577j_2 + 3363j_3, \\
 & BhC_3 = 99 + 577j_1 + 3363j_2 + 19601j_3, \\
 & \quad \quad \quad \vdots
 \end{aligned}$$

$$(7) \quad \begin{aligned} BhM_0 &= j_1 + 3j_2 + 7j_3, \\ BhM_1 &= 1 + 3j_1 + 7j_2 + 15j_3, \\ BhM_2 &= 3 + 7j_1 + 15j_2 + 31j_3, \\ BhM_3 &= 7 + 15j_1 + 31j_2 + 63j_3, \end{aligned}$$

$$(8) \quad \begin{aligned} &\vdots \\ BhH_0 &= 2 + 3j_1 + 5j_2 + 9j_3, \\ BhH_1 &= 3 + 5j_1 + 9j_2 + 17j_3, \\ BhH_2 &= 5 + 9j_1 + 17j_2 + 33j_3, \\ BhH_3 &= 9 + 17j_1 + 33j_2 + 65j_3, \end{aligned}$$

The following recurrence relations concerning the numbers BhB_n , BhC_n , BhM_n , BhH_n were proved in [5, 6].

Theorem 1.1 ([6]). *Let $n \geq 2$ be an integer. Then*

- (i) $BhB_n = 6BhB_{n-1} - BhB_{n-2}$,
- (ii) $BhC_n = 6BhC_{n-1} - BhC_{n-2}$,

where BhB_0 , BhB_1 , BhC_0 , BhC_1 are given by (5), (6), respectively.

Theorem 1.2 ([5]). *Let $n \geq 2$ be an integer. Then*

- (i) $BhM_n = 3BhM_{n-1} - 2BhM_{n-2}$,
- (ii) $BhH_n = 3BhH_{n-1} - 2BhH_{n-2}$,

where BhM_0 , BhM_1 , BhH_0 , BhH_1 are given by (7), (8), respectively.

For details concerning numbers BhB_n , BhC_n , BhM_n , BhH_n , including, among others, results about generating function and Binet's formulas see [5, 6].

In [1], Bednarz and Szynal-Liana proved relations between bihyperbolic numbers of the Fibonacci type and parameters of some special types of triangular tables and matrices. Following their research, in this paper we will express bihyperbolic numbers defined by formulas (1)–(4) as paraderminants and parapermanents of triangular matrices and as determinants and permanents of matrices. Before we do it, let us remind some facts about triangular matrices, paraderminants and parapermanents.

2. Triangular matrices, paraderminants and parapermanents.

An array of numbers from some field K of the form

$$A_n = \begin{bmatrix} a_{11} & & & & & \\ a_{21} & a_{22} & & & & \\ \vdots & \vdots & \ddots & & & \\ a_{n-1,1} & a_{n-1,2} & \cdots & a_{n-1,n-1} & & \\ a_{n1} & a_{n2} & \cdots & a_{n,n-1} & a_{nn} & \end{bmatrix}_{n \times n}$$

is known as a triangular matrix of order n .

It is important to acknowledge (see [24]) that a triangular matrix defined above is not a matrix in the classical sense since it is not a rectangular table of numbers.

Triangular matrices and special parameters connected with them, specifically paraterminants and parapermanents, are used in many branches of mathematics, see for example [13, 27]. For more information concerning triangular matrices, see [12, 25, 26, 24]. We will cite the most essential results, which are related to this paper.

In [12], the following formulas were given. Let A_n be a triangular matrix and by $\{a_{ij}\}$ let us denote the following expression

$$\{a_{ij}\} = \prod_{k=j}^i a_{ik}.$$

Then the paraterminant $\text{ddet}(A_n)$ and parapermanent $\text{pper}(A_n)$ of A_n are

$$\text{ddet}(A_n) = \sum_{r=1}^n \sum_{p_1+\dots+p_r=n} (-1)^{n-r} \prod_{s=1}^r \{a_{p_1+\dots+p_s, p_1+\dots+p_{s-1}+1}\}$$

and

$$\text{pper}(A_n) = \sum_{r=1}^n \sum_{p_1+\dots+p_r=n} \prod_{s=1}^r \{a_{p_1+\dots+p_s, p_1+\dots+p_{s-1}+1}\},$$

respectively, where summations are over the set of positive integer solutions of the equality $p_1 + \dots + p_r = n$.

For $n \geq 1$ we can decompose the paraterminant and the parapermanent by elements of the last row in the following way (see [12, 24]):

$$(9) \quad \text{ddet}(A_n) = \sum_{s=1}^n (-1)^{n-s} \{a_{ns}\} \text{ddet}(A_{s-1}),$$

$$(10) \quad \text{pper}(A_n) = \sum_{s=1}^n \{a_{ns}\} \text{pper}(A_{s-1}),$$

where $\text{ddet}(A_0) = 1$, $\text{pper}(A_0) = 1$.

In [28], Zatorsky and Lishchynskyy proved a relation between a paraterminant of a triangular matrix and a determinant of a special classical matrix, which is almost triangular, known as lower Hessenberg matrix. This

is the following relation:

$$(11) \quad \text{ddet}(A_n) = \det \begin{bmatrix} \{a_{11}\} & 1 & 0 & \cdots & 0 & 0 \\ \{a_{21}\} & \{a_{22}\} & 1 & \cdots & 0 & 0 \\ \{a_{31}\} & \{a_{32}\} & \{a_{33}\} & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \{a_{n-1,1}\} & \{a_{n-1,2}\} & \{a_{n-1,3}\} & \cdots & \{a_{n-1,n-1}\} & 1 \\ \{a_{n1}\} & \{a_{n2}\} & \{a_{n3}\} & \cdots & \{a_{n,n-1}\} & \{a_{nn}\} \end{bmatrix}.$$

Moreover, there exists a similar connection between parapermanent of a triangular matrix and a permanent of a lower Hessenberg matrix:

$$(12) \quad \text{pper}(A_n) = \text{per} \begin{bmatrix} \{a_{11}\} & 1 & 0 & \cdots & 0 & 0 \\ \{a_{21}\} & \{a_{22}\} & 1 & \cdots & 0 & 0 \\ \{a_{31}\} & \{a_{32}\} & \{a_{33}\} & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \{a_{n-1,1}\} & \{a_{n-1,2}\} & \{a_{n-1,3}\} & \cdots & \{a_{n-1,n-1}\} & 1 \\ \{a_{n1}\} & \{a_{n2}\} & \{a_{n3}\} & \cdots & \{a_{n,n-1}\} & \{a_{nn}\} \end{bmatrix}.$$

3. Main results. We are ready to present our results concerning non-trivial connections between special bihyperbolic sequences described in the introduction and paraderminants of triangular matrices.

Theorem 3.1. *Let $n \geq 0$ be an integer and let*

$$A_{n+1} = \begin{bmatrix} j_1 + 6j_2 + 35j_3 & & & & & & \\ -\frac{1}{6} + \frac{1}{6}j_2 + j_3 & 6 & & & & & \\ 0 & \frac{1}{6} & 6 & & & & \\ 0 & 0 & \frac{1}{6} & 6 & & & \\ \vdots & \vdots & \ddots & \ddots & \ddots & & \\ 0 & 0 & 0 & 0 & \frac{1}{6} & 6 & \end{bmatrix}_{(n+1) \times (n+1)}.$$

Then $BhB_n = \text{ddet}(A_{n+1})$.

Proof. (By induction with respect to n .)

If $n = 0$, then $\text{ddet}(A_1) = j_1 + 6j_2 + 35j_3 = BhB_0$.

If $n = 1$, then

$$\begin{aligned}
\text{ddet}(A_2) &= \sum_{s=1}^2 (-1)^{2-s} \{a_{2s}\} \text{ddet}(A_{s-1}) \\
&= (-1)^1 \cdot \{a_{21}\} \text{ddet}(A_0) + (-1)^0 \cdot \{a_{22}\} \text{ddet}(A_1) \\
&= (-1) \cdot \prod_{k=1}^2 a_{2k} \cdot \text{ddet}(A_0) + 1 \cdot \prod_{k=2}^2 a_{2k} \cdot \text{ddet}(A_1) \\
&= (-1) \cdot a_{21} \cdot a_{22} \cdot \text{ddet}(A_0) + 1 \cdot a_{22} \cdot \text{ddet}(A_1) \\
&= (-1) \cdot \left(-\frac{1}{6} + \frac{1}{6}j_2 + j_3 \right) \cdot 6 \cdot 1 + 1 \cdot 6 \cdot (j_1 + 6j_2 + 35j_3) \\
&= 1 - j_2 - 6j_3 + 6j_1 + 36j_2 + 210j_3 \\
&= 1 + 6j_1 + 35j_2 + 204j_3 = BhB_1.
\end{aligned}$$

Let us assume that for some integer $n \geq 0$ we have $BhB_n = \text{ddet}(A_{n+1})$ and $BhB_{n+1} = \text{ddet}(A_{n+2})$. We will show, that this assumption implies $BhB_{n+2} = \text{ddet}(A_{n+3})$. Using the formula (9), we obtain

$$\begin{aligned}
\text{ddet}(A_{n+3}) &= \sum_{s=1}^{n+3} (-1)^{n+3-s} \{a_{n+3,s}\} \text{ddet}(A_{s-1}) \\
&= (-1)^{n+3-1} \cdot \{a_{n+3,1}\} \text{ddet}(A_{1-1}) \\
&\quad + \dots + (-1)^{n+3-(n+1)} \cdot \{a_{n+3,n+1}\} \text{ddet}(A_{n+1-1}) \\
&\quad + (-1)^{n+3-(n+2)} \cdot \{a_{n+3,n+2}\} \text{ddet}(A_{n+2-1}) \\
&\quad + (-1)^{n+3-(n+3)} \cdot \{a_{n+3,n+3}\} \text{ddet}(A_{n+3-1}) \\
&= (-1)^{n+2} \cdot a_{n+3,1} \cdot a_{n+3,2} \cdot \dots \cdot a_{n+3,n+3} \cdot \text{ddet}(A_0) \\
&\quad + \dots + 1 \cdot a_{n+3,n+1} \cdot a_{n+3,n+2} \cdot \dots \cdot a_{n+3,n+3} \cdot \text{ddet}(A_n) \\
&\quad + (-1) \cdot a_{n+3,n+2} \cdot a_{n+3,n+3} \cdot \text{ddet}(A_{n+1}) \\
&\quad + 1 \cdot a_{n+3,n+3} \cdot \text{ddet}(A_{n+2}) \\
&= 0 + \dots + 0 + (-1) \cdot \frac{1}{6} \cdot 6 \cdot \text{ddet}(A_{n+1}) + 1 \cdot 6 \cdot \text{ddet}(A_{n+2}) \\
&= -\text{ddet}(A_{n+1}) + 6 \text{ddet}(A_{n+2}) \\
&= -BhB_n + 6BhB_{n+1} = BhB_{n+2},
\end{aligned}$$

which ends the proof. \square

By similar calculations we can prove analogous results for expressing BhC_n, BhM_n, BhH_n as paraderminants.

Proof. (By induction with respect to n .)

If $n = 0$, then $\text{pper}(A_1) = j_1 + 3j_2 + 7j_3 = BhM_0$.

If $n = 1$, then

$$\begin{aligned}
 \text{pper}(A_2) &= \sum_{s=1}^2 \{a_{2s}\} \text{pper}(A_{s-1}) = \{a_{21}\} \text{pper}(A_0) + \{a_{22}\} \text{pper}(A_1) \\
 &= \prod_{k=1}^2 a_{2k} \cdot \text{pper}(A_0) + \prod_{k=2}^2 a_{2k} \cdot \text{pper}(A_1) \\
 &= a_{21} \cdot a_{22} \cdot \text{pper}(A_0) + a_{22} \cdot \text{pper}(A_1) \\
 &= \left(\frac{1}{3} - \frac{2}{3}j_2 - 2j_3 \right) \cdot 3 \cdot 1 + 3 \cdot (j_1 + 3j_2 + 7j_3) \\
 &= 1 - 2j_2 - 6j_3 + 3j_1 + 9j_2 + 21j_3 \\
 &= 1 + 3j_1 + 7j_2 + 15j_3 = BhM_1.
 \end{aligned}$$

Let us assume that for some integer $n \geq 0$ we have $BhM_n = \text{pper}(A_{n+1})$ and $BhM_{n+1} = \text{pper}(A_{n+2})$. We will show, that this assumption implies $BhM_{n+2} = \text{pper}(A_{n+3})$. Applying the formula (10), we get

$$\begin{aligned}
 \text{pper}(A_{n+3}) &= \sum_{s=1}^{n+3} \{a_{n+3,s}\} \text{pper}(A_{s-1}) \\
 &= \{a_{n+3,1}\} \text{pper}(A_{1-1}) + \dots + \{a_{n+3,n+1}\} \text{pper}(A_{n+1-1}) \\
 &\quad + \{a_{n+3,n+2}\} \text{pper}(A_{n+2-1}) + \{a_{n+3,n+3}\} \text{pper}(A_{n+3-1}) \\
 &= a_{n+3,1} \cdot a_{n+3,2} \cdot \dots \cdot a_{n+3,n+3} \cdot \text{pper}(A_0) \\
 &\quad + \dots + a_{n+3,n+1} \cdot a_{n+3,n+2} \cdot \dots \cdot a_{n+3,n+3} \cdot \text{pper}(A_n) \\
 &\quad + a_{n+3,n+2} \cdot a_{n+3,n+3} \cdot \text{pper}(A_{n+1}) + a_{n+3,n+3} \cdot \text{pper}(A_{n+2}) \\
 &= 0 + \dots + 0 + \left(-\frac{2}{3} \right) \cdot 3 \cdot \text{pper}(A_{n+1}) + 3 \cdot \text{pper}(A_{n+2}) \\
 &= -2 \text{pper}(A_{n+1}) + 3 \text{pper}(A_{n+2}) \\
 &= -2BhM_n + 3BhM_{n+1} = BhM_{n+2},
 \end{aligned}$$

which end the proof. □

Theorem 3.6. *Let $n \geq 0$ be an integer and let*

$$A_{n+1} = \begin{bmatrix} j_1 + 6j_2 + 35j_3 & & & & & \\ \frac{1}{6} - \frac{1}{6}j_2 - j_3 & 6 & & & & \\ 0 & -\frac{1}{6} & 6 & & & \\ 0 & 0 & -\frac{1}{6} & 6 & & \\ \vdots & \vdots & \ddots & \ddots & \ddots & \\ 0 & 0 & 0 & 0 & -\frac{1}{6} & 6 \end{bmatrix}_{(n+1) \times (n+1)}.$$

Then $BhB_n = \text{pper}(A_{n+1})$.

$$BhM_n = \det \begin{bmatrix} j_1 + 3j_2 + 7j_3 & 1 & 0 & 0 & 0 & 0 \\ -1 + 2j_2 + 6j_3 & 3 & 1 & 0 & 0 & 0 \\ 0 & 2 & 3 & 1 & 0 & 0 \\ 0 & 0 & 2 & 3 & 1 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & 2 & 3 \end{bmatrix}_{(n+1) \times (n+1)},$$

$$BhH_n = \det \begin{bmatrix} 2 + 3j_1 + 5j_2 + 9j_3 & 1 & 0 & 0 & 0 & 0 \\ 3 + 4j_1 + 6j_2 + 10j_3 & 3 & 1 & 0 & 0 & 0 \\ 0 & 2 & 3 & 1 & 0 & 0 \\ 0 & 0 & 2 & 3 & 1 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & 2 & 3 \end{bmatrix}_{(n+1) \times (n+1)}.$$

Corollary 3.10. *Let $n \geq 0$ be an integer. Then*

$$BhB_n = \text{per} \begin{bmatrix} j_1 + 6j_2 + 35j_3 & 1 & 0 & \cdots & 0 & 0 \\ 1 - j_2 - 6j_3 & 6 & 1 & \cdots & 0 & 0 \\ 0 & -1 & 6 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & -1 & 6 & 1 \\ 0 & 0 & 0 & \cdots & -1 & 6 \end{bmatrix}_{(n+1) \times (n+1)},$$

$$BhC_n = \text{per} \begin{bmatrix} 1 + 3j_1 + 17j_2 + 99j_3 & 1 & 0 & \cdots & 0 & 0 \\ -3 - j_1 - 3j_2 - 17j_3 & 6 & 1 & \cdots & 0 & 0 \\ 0 & -1 & 6 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & -1 & 6 & 1 \\ 0 & 0 & 0 & \cdots & -1 & 6 \end{bmatrix}_{(n+1) \times (n+1)},$$

$$BhM_n = \text{per} \begin{bmatrix} j_1 + 3j_2 + 7j_3 & 1 & 0 & 0 & 0 & 0 \\ 1 - 2j_2 - 6j_3 & 3 & 1 & 0 & 0 & 0 \\ 0 & -2 & 3 & 1 & 0 & 0 \\ 0 & 0 & -2 & 3 & 1 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & -2 & 3 \end{bmatrix}_{(n+1) \times (n+1)},$$

$$BhH_n = \text{per} \begin{bmatrix} 2 + 3j_1 + 5j_2 + 9j_3 & 1 & 0 & 0 & 0 & 0 \\ -3 - 4j_1 - 6j_2 - 10j_3 & 3 & 1 & 0 & 0 & 0 \\ 0 & -2 & 3 & 1 & 0 & 0 \\ 0 & 0 & -2 & 3 & 1 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & -2 & 3 \end{bmatrix}_{(n+1) \times (n+1)}.$$

4. Examples. In this subsection we show how Corollary 3.10 and Theorem 3.1 work in specific cases, namely we express the number BhM_3 as a permanent of a matrix and the number BhB_3 as a paraderminant of a triangular matrix. In both cases, the permanent and the paraderminant are decomposed by elements of the last row.

$$\begin{aligned}
& \text{per} \begin{bmatrix} j_1 + 3j_2 + 7j_3 & 1 & 0 & 0 \\ 1 - 2j_2 - 6j_3 & 3 & 1 & 0 \\ 0 & -2 & 3 & 1 \\ 0 & 0 & -2 & 3 \end{bmatrix}_{4 \times 4} \\
&= 0 \cdot \text{per} \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ -2 & 3 & 1 \end{bmatrix}_{3 \times 3} + 0 \cdot \text{per} \begin{bmatrix} j_1 + 3j_2 + 7j_3 & 0 & 0 \\ 1 - 2j_2 - 6j_3 & 1 & 0 \\ 0 & 3 & 1 \end{bmatrix}_{3 \times 3} \\
&= (-2) \cdot \text{per} \begin{bmatrix} j_1 + 3j_2 + 7j_3 & 1 & 0 \\ 1 - 2j_2 - 6j_3 & 3 & 0 \\ 0 & -2 & 1 \end{bmatrix}_{3 \times 3} + 3 \cdot \text{per} \begin{bmatrix} j_1 + 3j_2 + 7j_3 & 1 & 0 \\ 1 - 2j_2 - 6j_3 & 3 & 1 \\ 0 & -2 & 3 \end{bmatrix}_{3 \times 3} \\
&= (-2) \cdot ((j_1 + 3j_2 + 7j_3) \cdot 3 + (1 - 2j_2 - 6j_3)) \\
&\quad + 3 \cdot ((j_1 + 3j_2 + 7j_3) \cdot 9 - 2 \cdot (j_1 + 3j_2 + 7j_3) + 3 \cdot (1 - 2j_2 - 6j_3)) \\
&= (-2) \cdot (1 + 3j_1 + 7j_2 + 15j_3) + 3 \cdot (3 + 7j_1 + 15j_2 + 31j_3) \\
&= 7 + 15j_1 + 31j_2 + 63j_3 = BhM_3.
\end{aligned}$$

$$\begin{aligned}
& \text{ddet} \begin{bmatrix} j_1 + 6j_2 + 35j_3 & & & \\ -\frac{1}{6} + \frac{1}{6}j_2 + j_3 & 6 & & \\ 0 & \frac{1}{6} & 6 & \\ 0 & 0 & \frac{1}{6} & 6 \end{bmatrix}_{4 \times 4} \\
&= (-1)^3 \cdot \left(0 \cdot 0 \cdot \frac{1}{6} \cdot 6\right) \cdot 1 + (-1)^2 \cdot \left(0 \cdot \frac{1}{6} \cdot 6\right) \cdot \text{ddet} [j_1 + 6j_2 + 35j_3]_{1 \times 1} \\
&\quad + (-1)^1 \cdot \left(\frac{1}{6} \cdot 6\right) \cdot \text{ddet} \begin{bmatrix} j_1 + 6j_2 + 35j_3 & \\ -\frac{1}{6} + \frac{1}{6}j_2 + j_3 & 6 \end{bmatrix}_{2 \times 2} \\
&\quad + (-1)^0 \cdot 6 \cdot \text{ddet} \begin{bmatrix} j_1 + 6j_2 + 35j_3 & & \\ -\frac{1}{6} + \frac{1}{6}j_2 + j_3 & 6 & \\ 0 & \frac{1}{6} & 6 \end{bmatrix}_{3 \times 3} \\
&= 0 + 0 - \text{ddet} \begin{bmatrix} j_1 + 6j_2 + 35j_3 & \\ -\frac{1}{6} + \frac{1}{6}j_2 + j_3 & 6 \end{bmatrix}_{2 \times 2} + 6 \cdot \text{ddet} \begin{bmatrix} j_1 + 6j_2 + 35j_3 & & \\ -\frac{1}{6} + \frac{1}{6}j_2 + j_3 & 6 & \\ 0 & \frac{1}{6} & 6 \end{bmatrix}_{3 \times 3} \\
&= -\text{ddet} \begin{bmatrix} j_1 + 6j_2 + 35j_3 & \\ -\frac{1}{6} + \frac{1}{6}j_2 + j_3 & 6 \end{bmatrix}_{2 \times 2} + 6 \cdot (-1)^1 \cdot \left(\frac{1}{6} \cdot 6\right) \cdot \text{ddet} [j_1 + 6j_2 + 35j_3]_{1 \times 1} \\
&\quad + 6 \cdot (-1)^0 \cdot 6 \cdot \text{ddet} \begin{bmatrix} j_1 + 6j_2 + 35j_3 & & \\ -\frac{1}{6} + \frac{1}{6}j_2 + j_3 & 6 & \end{bmatrix}_{2 \times 2}
\end{aligned}$$

$$\begin{aligned}
&= 35 \cdot \text{ddet} \begin{bmatrix} j_1 + 6j_2 + 35j_3 & \\ -\frac{1}{6} + \frac{1}{6}j_2 + j_3 & 6 \end{bmatrix}_{2 \times 2} - 6 \cdot \text{ddet} [j_1 + 6j_2 + 35j_3]_{1 \times 1} \\
&= 35 \cdot \left((-1)^1 \cdot \left(-\frac{1}{6} + \frac{1}{6}j_2 + j_3 \right) \cdot 6 \cdot 1 + (-1)^0 \cdot 6 \cdot \text{ddet} [j_1 + 6j_2 + 35j_3]_{1 \times 1} \right) \\
&\quad + \left(-6 \cdot \text{ddet} [j_1 + 6j_2 + 35j_3]_{1 \times 1} \right) \\
&= 35 \cdot (1 - j_2 - 6j_3) + 204 \cdot \text{ddet} [j_1 + 6j_2 + 35j_3]_{1 \times 1} \\
&= 35 - 35j_2 - 210j_3 + 204 \cdot (j_1 + 6j_2 + 35j_3) \\
&= 35 + 204j_1 + 1189j_2 + 6930j_3 \\
&= BhB_3.
\end{aligned}$$

5. Conclusions. In our paper we have considered special bihyperbolic numbers given by linear recurrence of the second order. We have shown that these numbers can be expressed as certain parameters of specific triangular tables or matrices. The obtained results may be a starting point for the search of further relations between matrix theory and bihyperbolic numbers defined by linear or nonlinear recurrence equations of order $k \geq 3$, for example the well-known Padovan recurrence given by the formula $P(n) = P(n-2) + P(n-3)$, where $P(0) = P(1) = P(2) = 1$.

REFERENCES

- [1] Bednarz, P., Szynal-Liana, A., *Bihyperbolic numbers of the Fibonacci type and triangular matrices (tables)*, Azerb. J. Math. **14**(2) (2024), 79–87.
- [2] Behera, A., Panda, G. K., *On the square roots of triangular numbers*, Fibonacci Quart. **37**(2) (1999), 98–105.
- [3] Bilgin, M., Ersoy, S., *Algebraic properties of bihyperbolic numbers*, Adv. Appl. Clifford Algebr. **30**(1) (2020), 1–17.
- [4] Boussayoud, A., Chelgham, M., Boughaba, S., *On some identities and generating functions for Mersenne numbers and polynomials*, Turk. J. Anal. Numb. Theory **6**(3) (2018), 93–97.
- [5] Bród, D., Szynal-Liana, A., *On Mersenne numbers and their bihyperbolic generalizations*, Ann. Math. Sil. **39**(1) (2025), 130–142.
- [6] Bród, D., Szynal-Liana, A., Włoch, I., *On the combinatorial properties of bihyperbolic balancing numbers*, Tatra Mt. Math. Publ. **77** (2020), 27–38.
- [7] Bród, D., Szynal-Liana, A., Włoch, I., *Bihyperbolic numbers of the Fibonacci type and their idempotent representation*, Comment. Math. Univ. Carolin. **62**(4) (2021), 409–416.
- [8] Catarino, P., Campos, H., Vasco, P., *On some identities for balancing and cobalancing numbers*, Ann. Math. Inform. **45** (2015), 11–24.
- [9] Catarino, P., Campos, H., Vasco, P., *On the Mersenne sequence*, Ann. Math. Inform. **46** (2016), 37–53.
- [10] Cockle, J., *On certain functions resembling quaternions, and on a new imaginary in algebra*, Lond. Edinb. Dublin Philos. Mag. J. Sci. **33** (1848), 435–439.
- [11] Davala, R. K., Panda, G. K., *On sum and ratio formulas for Lucas-balancing numbers*, Palest. J. Math. **8**(2) (2019), 200–206.

- [12] Ganyushkin, O. G., Zatorsky, R. A., Lishchinskyy, I. I., *On paraderminants and parapermanents of triangular matrices*, Bull. Kyiv Univ. Ser. Phys. Math. **1** (2005), 35–41.
- [13] Goy, T., Zatorsky, R. A., *Infinite linear recurrence relation and superposition of linear recurrence equations*, J. Integer Seq. **20**(5) (2017), Article 17.5.3.
- [14] Gürses, N., İşbilir Z., *An extended framework for bihyperbolic generalized Tribonacci numbers*, Commun. Fac. Sci. Univ. Ank. Ser. A1. Math. Stat. **73**(3), (2024), 765–786.
- [15] Kantor, I. L., Solodovnikov, A. S., *Hypercomplex Numbers. An Elementary Introduction to Algebras*, 1st edn. Springer, New York, 1989.
- [16] Ochalik, P., Włoch, A., *On generalized Mersenne numbers, their interpretations and matrix generators*, Ann. Univ. Mariae Curie-Skłodowska Sectio A **72**(1) (2018), 69–76.
- [17] Olariu, S., *Complex Numbers in n Dimensions*, Elsevier, Amsterdam, 2002.
- [18] Panda, G. K., *Some fascinating properties of balancing numbers*, in: *Proceedings of The Eleventh International Conference on Fibonacci Numbers and Their Applications*, 185–189, Cong. Numerantium **194** (2009).
- [19] Pogorui, A. A., Rodriguez-Dagnino, R. M., Rodrigue-Said, R. D., *On the set of zeros of bihyperbolic polynomials*, Complex Var. Elliptic Equ. **53**(7) (2008), 685–690.
- [20] Rayaguru, S. G., Bravo, J. J., *Balancing and Lucas-balancing numbers which are concatenation of three repdigits*, Bol. Soc. Mat. Mex. **29**(3) (2023), Paper No. 57.
- [21] Rochon, D., Shapiro M., *On algebraic properties of bicomplex and hyperbolic numbers*, An. Univ. Oradea Fasc. Mat. **11** (2004), 71–110.
- [22] Saba, N., Boussayoud, A., Kanuri, K. V. V., *Mersenne Lucas numbers and complete homogeneous symmetric functions*, J. Math. Computer Sci. **24** (2022), 127–139.
- [23] Yaglom, I. M., *A Simple non-Euclidean Geometry and its Physical Basis*, Springer-Verlag, New York-Heidelberg, 1979.
- [24] Zatorsky, R. A., *Theory of paraderminants and its applications*, Algebra Discrete Math. **1** (2007), 108–137.
- [25] Zatorsky, R. A., *Introduction to the theory of triangular matrices (tables)*, in: I. I. Kyrchev (Ed.), *Advances in Linear Algebra Research*, Nova Science Publishers, New York, 2015, pp. 185–238.
- [26] Zatorsky, R. A., *On paraderminants and parapermanents of triangular matrices*, Mat. Stud. **17**(1) (2002), 3–17.
- [27] Zatorsky, R. A., Goy, T., *Parapermanents of triangular matrices and some general theorems on number sequences*, J. Integer Seq. **19** (2016), Article 16.2.2.
- [28] Zatorsky, R. A., Lishchynskyy, I. I., *On connection between determinants and paraderminants*, Mat. Stud. **25** (2006), 97–102.

Paweł Bednarz
Rzeszów University of Technology
The Faculty of Mathematics
and Applied Physics
Department of Discrete Mathematics
al. Powstańców Warszawy 12
35-959 Rzeszów
Poland
e-mail: pbednarz@prz.edu.pl

Anna Kosiorowska
Rzeszów University of Technology
The Faculty of Mathematics
and Applied Physics
Department of Discrete Mathematics
al. Powstańców Warszawy 12
35-959 Rzeszów
Poland
e-mail: a.kosiorowsk@prz.edu.pl

Adrian Michalski
Rzeszow University of Technology
The Faculty of Mathematics and Applied Physics
Department of Discrete Mathematics
al. Powstańców Warszawy 12
35-959 Rzeszów
Poland
e-mail: a.michalski@prz.edu.pl

Received April 9, 2025